arXiv:math/0302357v1 [math.CA] 28 Feb 2003

Quadratic Hermite–Padé approximation to the exponential function: a Riemann–Hilbert approach*

A. B. J. Kuijlaars, W. Van Assche,
Katholieke Universiteit Leuven, Belgium

and

F. Wielonsky
Université des Sciences et Technologies Lille 1, France, and
INRIA Sophia Antipolis, France

February 1, 2008

Abstract

We investigate the asymptotic behavior of the polynomials p, q, r of degrees n in type I Hermite-Padé approximation to the exponential function, defined by $p(z)e^{-z} + q(z) + r(z)e^{z} = \mathcal{O}(z^{3n+2})$ as $z \to 0$. These polynomials are characterized by a Riemann–Hilbert problem for a 3×3 matrix valued function. We use the Deift-Zhou steepest descent method for Riemann–Hilbert problems to obtain strong uniform asymptotics for the scaled polynomials $p(3nz)$, $q(3nz)$, and $r(3nz)$ in every domain in the complex plane. An important role is played by a three-sheeted Riemann surface and certain measures and functions derived from it. Our work complements recent results of Herbert Stahl.

Contents

1 Introduction **2**

2 Statement of results **6**
2.1 The Riemann surface 6
2.2 The measures μ_P, μ_Q, μ_R, and μ_E 9
2.3 The g-functions 11
2.4 Functions φ_P and φ_R 12
2.5 Strong asymptotics away from the zeros 12
2.6 Asymptotics near the curves Γ_P, Γ_Q, Γ_R and Γ_E 14
2.7 Asymptotics near the branch points 15
2.8 Convergence of the Hermite–Padé approximants 17

3 Geometry of the problem **18**
3.1 Proof of Propositions 2.1 and 2.2 18

---

[*] This work is supported by INTAS 2000-0272, projects G.0176.02 and G.0184.02 of FWO-Vlaanderen

4 Measures and functions associated with the Riemann surface 21
4.1 Properties of the measures μ_P, μ_R, and μ_E 21
4.2 Properties of g_P and g_R 23
4.3 Jump properties of g_P and g_R 24
4.4 Proof of Lemma 3.1 26
4.5 Proof of Lemma 2.7 26
4.6 The measure μ_Q 27
4.7 The function g_Q 28

5 The Riemann–Hilbert problem and the first two transformations 29
5.1 The Riemann–Hilbert problem 29
5.2 First transformation 31
5.3 Deformation of contours 32

6 Construction of parametrices and final transformation 36
6.1 Parametrix for the exterior region 36
6.2 Parametrices near the branch points 40
6.3 Third transformation 45

7 Proofs of the asymptotic formulas 46
7.1 Proof of Theorem 2.8 46
7.2 Proof of Theorem 2.5 49
7.3 Proof of Theorem 2.9 52
7.4 Proof of Theorem 2.10 53
7.5 Proof of Theorem 2.11 54
7.6 Proof of Corollary 2.12 55
7.7 Proof of Theorem 2.13 56

References 57

1 Introduction

The study of Hermite–Padé approximation for the exponential function was initiated by Hermite [19], in connection with his proof of the transcendency of e. A special case of this is the quadratic Hermite–Padé approximation of type I, where for any given three integers $n_1, n_2, n_3 \geq 0$, one asks for polynomials p_{n_1,n_2,n_3}, q_{n_1,n_2,n_3}, and r_{n_1,n_2,n_3} of degrees n_1, n_2, n_3, respectively, such that

$$p_{n_1,n_2,n_3}(z) + q_{n_1,n_2,n_3}(z)e^{z} + r_{n_1,n_2,n_3}(z)e^{2z} = \mathcal{O}(z^{n_1+n_2+n_3+2}), \qquad z \to 0. \tag{1.1}$$

These polynomials exist and are unique up to a common multiplicative factor. We put

$$e_{n_1,n_2,n_3}(z) = p_{n_1,n_2,n_3}(z)e^{-z} + q_{n_1,n_2,n_3}(z) + r_{n_1,n_2,n_3}(z)e^{z},$$

so that

$$e_{n_1,n_2,n_3}(z) = \mathcal{O}(z^{n_1+n_2+n_3+2}), \qquad z \to 0.$$

Putting the left-hand side of (1.1) equal to zero, we obtain the quadratic algebraic Hermite–Padé approximants

$$\frac{-q_{n_1,n_2,n_3}(z) \pm \sqrt{q_{n_1,n_2,n_3}(z)^2 - 4p_{n_1,n_2,n_3}(z)r_{n_1,n_2,n_3}(z)}}{2r_{n_1,n_2,n_3}(z)}$$

to e^z.

The analytic theory of the Hermite–Padé approximation to the exponential function was investigated, among others, by Mahler [26], Aptekarev [1], Chudnovsky [11], Borwein [10], Driver [16], Driver and Temme [17], and Wielonsky [41, 42]. The paper [27] by Nuttall deserves a special mention since it contains many deep ideas and conjectures about the asymptotics of Hermite–Padé approximants to general functions. Several surveys on the subject also exist, see De Bruin [9], Baker and Lubinsky [5], and Aptekarev and Stahl [2].

Very recently, Herbert Stahl [34, 35, 36] gave detailed asymptotic results for the scaled diagonal polynomials

$$P_n(z) = p_{n,n,n}(3nz), \qquad Q_n(z) = q_{n,n,n}(3nz), \qquad R_n(z) = r_{n,n,n}(3nz), \tag{1.2}$$

and for the remainder term

$$E_n(z) = P_n(z)e^{-3nz} + Q_n(z) + R_n(z)e^{3nz} \tag{1.3}$$

in the approximation. From these asymptotics, the limit distributions of the zeros of P_n, Q_n, R_n and E_n follow. The zeros accumulate on specific arcs in the complex plane. In Figure 1, the zeros of $p_{n,n,n}$, $q_{n,n,n}$, and $r_{n,n,n}$ have been plotted for the value $n = 60$. It shows their remarkable distribution, especially the zeros of $q_{n,n,n}$ distribute themselves in a very particular way. These results on the distribution of the zeros of Hermite–Padé approximants to exponentials may be seen as a continuation of the study initiated by Szegő in [38] concerning the distribution of the zeros of Taylor sections of the series for e^z, subsequently generalized by Saff and Varga in [31, 32, 33] to the zeros of the Padé approximants to e^z, see also [22, 40].

Stahl [34, 35, 36] obtained his results from a saddle point analysis for the explicit integral formulas for P_n, Q_n, and R_n, which are

$$P_n(z) = \frac{Ce^{3nz}}{2\pi i}\oint_{C_{-1}} \frac{e^{3nzw}dw}{[w(w^2-1)]^{n+1}}, \tag{1.4}$$

$$Q_n(z) = \frac{C}{2\pi i}\oint_{C_0} \frac{e^{3nzw}dw}{[w(w^2-1)]^{n+1}}, \tag{1.5}$$

$$R_n(z) = \frac{Ce^{-3nz}}{2\pi i}\oint_{C_1} \frac{e^{3nzw}dw}{[w(w^2-1)]^{n+1}}. \tag{1.6}$$

Here C_j is a closed contour in the complex plane encircling j in the positive direction, which does not encircle the other points in $\{-1, 0, 1\}$. The number C in (1.4)–(1.6) is a normalization constant. Stahl used $C = n!2^{n+1}(3n)^{-n}$ in order to make P_n a monic polynomial. For our analysis below, we found it convenient to make Q_n monic, and we choose

$$C = \frac{n!(-1)^{n+1}}{(3n)^n}. \tag{1.7}$$



Figure 1: Zeros of the diagonal quadratic Hermite–Padé polynomials $p_{60,60,60}$ (the diamonds on the left) $q_{60,60,60}$ (the circles in the middle), and $r_{60,60,60}$ (the boxes on the right)

In this paper we propose a different approach to the asymptotic analysis. Our approach is based on a Riemann–Hilbert formulation for the polynomials P_n, Q_n, and R_n, combined with a steepest descent analysis for Riemann–Hilbert problems. This technique originated with Deift and Zhou [15] and was applied to the asymptotics of orthogonal polynomials by Deift et al. [12, 13, 14]. See also [7, 8, 4, 20, 21, 22, 23, 24] for recent developments. The Riemann–Hilbert problem for multiple orthogonal polynomials was given by Van Assche et al. [39]. In the present situation it gives rise to a Riemann–Hilbert problem for a 3×3-matrix valued function. As far as we are aware, this is the first time that the steepest descent method for Riemann–Hilbert problems is applied to a 3×3 problem.

The Riemann–Hilbert problem is to find a 3×3 matrix valued function $Y : \mathbb{C} \setminus \Gamma \to \mathbb{C}^{3\times 3}$ where Γ is a closed contour in the complex plane encircling the origin once in the positive direction, such that

1. Y is analytic in $\mathbb{C} \setminus \Gamma$.

2. Y satisfies the jump condition

$$Y_+(z) = Y_-(z) \begin{pmatrix} 1 & z^{-3n-2}e^{-3nz} & 0 \\ 0 & 1 & 0 \\ 0 & z^{-3n-2}e^{3nz} & 1 \end{pmatrix}, \qquad z \in \Gamma, \tag{1.8}$$

where $Y_+(z)$ and $Y_-(z)$ denote the limiting values of $Y(z')$ as z' approaches $z \in \Gamma$ from the inside and outside of Γ, respectively.

3. For large z

$$Y(z) = \left(I + \mathcal{O}\left(\frac{1}{z}\right)\right) \begin{pmatrix} z^{n+1} & 0 & 0 \\ 0 & z^{-2n-2} & 0 \\ 0 & 0 & z^{n+1} \end{pmatrix}, \qquad z \to \infty. \tag{1.9}$$

We will show in Section 5.1 that the Riemann–Hilbert problem has a unique solution and that $Y_{21}(z) = P_n(z)$, $Y_{22}(z) = z^{-3n-2}Q_n(z)$ for z outside Γ, $Y_{22}(z) = z^{-3n-2}E_n(z)$ for z inside Γ, and $Y_{23}(z) = R_n(z)$.

The steepest descent analysis consists of a number of transformations. To make the transformations work, we make heavy use of the works of Stahl [34, 35, 36]. A crucial role is played by the Riemann surface defined by

$$z = \frac{w^2 - \frac{1}{3}}{w(w^2 - 1)}, \tag{1.10}$$

which is considered as a three sheeted surface with cuts along two arcs Γ_P and Γ_R. The jumps of the different inverse mappings of (1.10) across the arcs determine probability measures μ_P and μ_R supported on Γ_P and Γ_R, respectively. These measures turn out to be limiting distributions of the normalized zero counting measures of the polynomials P_n and R_n, respectively.

We choose the closed contour Γ in the Riemann–Hilbert problem for Y so that it contains the cuts Γ_P and Γ_R. The measures μ_P and μ_R and their g-transforms

$$g_P(z) = \int \log(z - s) d\mu_P(s), \qquad g_R(z) = \int \log(z - s) d\mu_R(s)$$

are used to make the first transformation of the Riemann–Hilbert problem, which has the effect of normalizing the problem at infinity. Then we follow the general scheme, as presented in [14] or [12], for the asymptotic analysis of Riemann–Hilbert problems. It leads to a final Riemann–Hilbert problem whose solution has an explicit asymptotic behavior for $n \to \infty$, see Theorem 6.4 in Section 6.3. Tracing our steps back to the original Riemann–Hilbert problem, we obtain strong asymptotic formulas for the Hermite–Padé approximants in every region of the complex plane, including the sets where the zeros are and their endpoints.

In Section 2 we state the asymptotic results for the polynomials P_n, Q_n, and R_n, and for the remainder E_n, as well as for the quadratic Hermite–Padé approximants. The asymptotic results make use of the functions obtained from the Riemann surface. The Riemann surface and the measures and functions derived from it are also described in Section 2. In Section 3 we prove the statements about the Riemann surface and other geometrical objects involved in the problem. To prepare for the transformations of the Riemann–Hilbert problem we need a large number of relations between the various functions involved, such as the inverse mappings of (1.10) and the functions g_P and g_R. These properties are established in terms in Section 4. Sections 5 and 6 contain the transformations of the Riemann–Hilbert problem and all asserted asymptotic results are finally proven in Section 7.

2 Statement of results

2.1 The Riemann surface

In order to state our results we first introduce an appropriate Riemann surface. The Riemann surface is chosen so that it parameterizes the critical points of the function

$$w \mapsto 3zw - \log\left[w(w^2-1)\right]. \tag{2.1}$$

Note that the integrals in formulas (1.4)–(1.6) have the form

$$\oint_{C_j} \frac{1}{w(w^2-1)} e^{n\left(3zw - \log\left[w(w^2-1)\right]\right)} dw \tag{2.2}$$

and that by the classical saddle point analysis for the asymptotic evaluation of integrals, the main contribution to the integral (2.2) comes from a critical point of (2.1). So we define $\mathcal{R}$ as the Riemann surface for the function

$$z = z(w) = \frac{1}{3}\left(\frac{1}{w} + \frac{1}{w-1} + \frac{1}{w+1}\right) = \frac{w^2 - \frac{1}{3}}{w(w^2-1)}. \tag{2.3}$$

Note that we obtain (2.3) if we set the derivative of (2.1) equal to zero and solve for z.



Figure 2: The Riemann surface $\mathcal{R}$

The rational function (2.3) has three inverse mappings. These are the three solutions of the cubic equation

$$zw^3 - w^2 - zw + \frac{1}{3} = 0. \tag{2.4}$$

The Riemann surface $\mathcal{R}$ consists of three sheets $\mathcal{R}_P$, $\mathcal{R}_Q$, and $\mathcal{R}_R$ as shown in Figure 2, see Proposition 2.1. The bijective mapping $\psi : \mathcal{R} \to \overline{\mathbb{C}}$ is the inverse of (2.3). We denote its restriction to the three sheets by ψ_P, ψ_Q, and ψ_R, respectively. So $\psi_P(z)$, $\psi_Q(z)$, and $\psi_R(z)$

are the three solutions of (2.4). Typically we will identify the three sheets with copies of the complex plane, and so ψ_P, ψ_Q, and ψ_R are defined on $\mathbb{C}$ with appropriate cuts. The sheets $\mathcal{R}_P$ and $\mathcal{R}_Q$ are glued together along a cut Γ_P connecting two branch points z_1 and z_2, and the sheets $\mathcal{R}_Q$ and $\mathcal{R}_R$ are glued together along a cut Γ_R connecting the other two branch points z_3 and z_4.

The Riemann surface has four branch points $z_1 = z(w_1)$, $z_2 = z(w_2)$, $z_3 = z(w_3)$, $z_4 = z(w_4)$, which are related to the points w_1, w_2, w_3, w_4 for which $z'(w) = 0$. Simple calculations give

$$w_k = 3^{-1/4}\omega_8^{-2k-1}, \qquad k = 1,2,3,4, \tag{2.5}$$

where $\omega_8 = e^{2\pi i/8}$ is the primitive 8th root of unity. The corresponding values of $z_k = z(w_k)$ are

$$z_1 = 3^{-1/4}\omega_{24}^{7}, \quad z_2 = 3^{-1/4}\omega_{24}^{17}, \quad z_3 = 3^{-1/4}\omega_{24}^{19}, \quad z_4 = 3^{-1/4}\omega_{24}^{5}, \tag{2.6}$$

where $\omega_{24} = e^{2\pi i/24}$ is the primitive 24th root of unity. The precise sheet structure of $\mathcal{R}$ is given in the following proposition.

Proposition 2.1. *There is an analytic curve Γ_P from z_1 to z_2 lying in the left half-plane, and an analytic curve Γ_R from z_3 to z_4 lying in the right half-plane, such that the following hold.*

(a) *Three inverse mappings ψ_P, ψ_Q, and ψ_R of* (2.3) *exist so that ψ_P is defined and analytic on $\mathbb{C}\setminus\Gamma_P$, ψ_Q is defined and analytic on $\mathbb{C}\setminus(\Gamma_P\cup\Gamma_R)$, and ψ_R is defined and analytic on $\mathbb{C}\setminus\Gamma_R$.*

(b) *At infinity, we have the values $\psi_P(\infty) = -1$, $\psi_Q(\infty) = 0$, and $\psi_R(\infty) = 1$.*

(c) *For $z \in \Gamma_P$, we have*

$$\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)_+(s)ds \in \mathbb{R}, \tag{2.7}$$

with integration along the +side of Γ_P. [This is the side of Γ_P that is on the left while going from z_1 to z_2 along Γ_P.]

(d) *For $z \in \Gamma_R$, we have*

$$\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q - \psi_R)_+(s)\,ds \in \mathbb{R}, \tag{2.8}$$

with integration along the +side of Γ_R.

Because of symmetry, Γ_R is the mirror image of Γ_P under reflection with respect to the imaginary axis. We take the sheet structure on the Riemann surface $\mathcal{R}$ as in Figure 2. The functions ψ_P, ψ_Q, ψ_R are defined on the P, Q, and R sheet of $\mathcal{R}$, respectively. Together they constitute a conformal map from $\mathcal{R}$ onto the Riemann sphere. The images of the different sheets are shown in Figure 3.

Other curves of interest to our problem are defined by the property that either $\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)(s)ds$ or $\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q - \psi_R)(s)ds$ is real. These curves are described by the following proposition, see Figure 4.

Proposition 2.2. (a) *There are four analytic curves where $\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)(s)\,ds$ is real. One of them is Γ_P, a second one joins z_2 with z_1 and intersects the positive real axis. We call this curve Γ_P^*. The other two curves are unbounded and lie in the left half-plane. They join z_1 and z_2 with infinity, and we call them $\Gamma_{E,1}$ and $\Gamma_{E,2}$, respectively.*



Figure 3: ψ-image of the Riemann surface $\mathcal{R}$

(b) *There are four analytic curves where* $\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q-\psi_R)(s)\,ds$ *is real. One of them is* Γ_R*, a second one joins* z_4 *with* z_3 *and intersects the negative real axis. We call this curve* Γ_R^**. The other two curves are unbounded and lie in the right half-plane. They join* z_3 *and* z_4 *with infinity and we call them* $\Gamma_{E,3}$ *and* $\Gamma_{E,4}$*, respectively.*

(c) *The curves* Γ_P^* *and* Γ_R *do not intersect.*

(d) *The curves* Γ_R^* *and* Γ_P *do not intersect.*

(e) *The curves* Γ_P^* *and* Γ_R^* *intersect in the points* $\pm iy^*$ *on the imaginary axis. The value of* y^* *is approximately* $y^* = 0.621391\cdots$.

By symmetry, Γ_P^* and Γ_R^* are mirror images with respect to the imaginary axis. Similarly, $\Gamma_{E,1}$ and $\Gamma_{E,4}$ are mirror images of each other, as well as $\Gamma_{E,2}$ and $\Gamma_{E,3}$. All contours are oriented as shown in Figure 4. The orientation induces a +side and a −side on each contour, where the +side is on the left and the −side on the right while traversing the contour according to its orientation. Propositions 2.1 and 2.2 are proved in Section 3.

Now we can also define the contours Γ_Q and Γ_E. We define

$$\Gamma_E = \bigcup_{k=1}^{4} \Gamma_{E,k}. \tag{2.9}$$

where $\Gamma_{E,k}$, $k = 1,2,3,4$ are as in Proposition 2.2, and

$$\Gamma_Q = [-iy^*, iy^*] \cup (\Gamma_P^* \cap \{\operatorname{Re} z < 0\}) \cup (\Gamma_R^* \cap \{\operatorname{Re} z > 0\}), \tag{2.10}$$

where Γ_P^*, Γ_R^* and iy^* are as in Proposition 2.2. We choose an orientation on Γ_Q by orienting the interval $[-iy^*, iy^*]$ from $-iy^*$ to iy^* and by orienting Γ_P^* from z_2 to z_1 and Γ_R^* from z_4 to z_3, see Figure 5.



Figure 4: Curves for which either $\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q-\psi_P)(s)\,ds$ or $\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q-\psi_R)(s)\,ds$ is real.

The contours Γ_P, Γ_Q, Γ_R, and Γ_E divide the complex plane into six domains. We denote the unbounded domains by $D_{\infty,P}$, $D_{\infty,R}$, $D_{\infty,U}$ and $D_{\infty,L}$, as shown in Figure 5. The bounded domains are denoted by D_P and D_R, where D_P is in the left half-plane, and D_R is in the right half-plane, see also Figure 5. We also put

$$D_\infty = (D_{\infty,P}\cup D_{\infty,R}\cup D_{\infty,U}\cup D_{\infty,L})\cup(\Gamma_E\setminus\{z_1,z_2,z_3,z_4\}). \tag{2.11}$$

This is the unbounded domain bounded by Γ_P, Γ_Q and Γ_R. Of further interest is the bounded domain D_P^* bounded by Γ_P and Γ_P^*, see Figure 6, and its mirror image D_R^* with respect to the imaginary axis.

This concludes the description of the Riemann surface.

2.2 The measures μ_P, μ_Q, μ_R, and μ_E

We now define a measure on each of the curves Γ_P, Γ_Q, Γ_R, and Γ_E. The complex line element ds is defined according to the orientation of these curves given in Figure 5.

Definition 2.3. We define a measure μ_P on Γ_P by

$$d\mu_P(s)=\frac{3}{2\pi i}(\psi_Q-\psi_P)_+(s)\,ds \qquad s\in\Gamma_P, \tag{2.12}$$

a measure μ_R on Γ_R by

$$d\mu_R(s)=\frac{3}{2\pi i}(\psi_Q-\psi_R)_+(s)\,ds \qquad s\in\Gamma_R, \tag{2.13}$$



Figure 5: Curves Γ_P, Γ_Q, Γ_R, and Γ_E, and domains D_P, D_R, $D_{\infty,P}$,$D_{\infty,R}$,$D_{\infty,U}$, and $D_{\infty,L}$. Note that Γ_Q does not only consist of the interval $[-iy^*, iy^*]$ but also of the small arcs from z_2 via $-iy^*$ to z_3 and from z_4 via iy^* to z_1. The domain D_∞ is the union of Γ_E with the unbounded domains $D_{\infty,P}$, $D_{\infty,R}$, $D_{\infty,U}$, and $D_{\infty,L}$.

and a measure μ_Q on Γ_Q by

$$d\mu_Q(s) = \begin{cases} \frac{3}{2\pi i}(\psi_Q - \psi_P)(s)\,ds & s \in \Gamma_P^* \cap \{\operatorname{Re} z < 0\}, \\ \frac{3}{2\pi i}(\psi_Q - \psi_R)(s)\,ds & s \in \Gamma_R^* \cap \{\operatorname{Re} z > 0\}, \\ \frac{3}{2\pi i}(\psi_R - \psi_P)(s)\,ds & s \in [-iy^*, iy^*]. \end{cases} \tag{2.14}$$

The measure μ_E on Γ_E is defined as

$$d\mu_E(s) = \begin{cases} \frac{3}{2\pi i}(\psi_Q - \psi_P)(s)\,ds & s \in \Gamma_{E,1} \cup \Gamma_{E,2}, \\ \frac{3}{2\pi i}(\psi_Q - \psi_R)(s)\,ds & s \in \Gamma_{E,3} \cup \Gamma_{E,4}. \end{cases} \tag{2.15}$$

A priori these are complex measures. Our first result is that μ_P, μ_Q, and μ_R are in fact probability measures and μ_E is a positive measure.

Theorem 2.4. *We have that μ_P is a probability measure on Γ_P, μ_Q is a probability measure on Γ_Q and μ_R is a probability measure on Γ_R. The measure μ_E is a positive measure on Γ_E.*

Theorem 2.4 follows from Propositions 4.2 and 4.8.

The relevance of these measures is shown by the following theorem. For every polynomial p of exact degree n, we denote by ν_p the normalized zero counting measure. Thus

$$\nu_p = \frac{1}{n} \sum_{p(z)=0} \delta_z$$

where each zero is counted according to its multiplicity. We also define a zero counting measure for the remainder function E_n, namely

$$\nu_{E_n} = \frac{1}{n} \sum_{\substack{E_n(z)=0 \\ z\neq 0}} \delta_z,$$

where the normalization by n now corresponds to the degree of approximation and the $3n+2$ interpolatory zeros of E_n at the origin have been excluded.

Theorem 2.5. *We have*

$$\nu_{P_n} \xrightarrow{*} \mu_P, \qquad \nu_{Q_n} \xrightarrow{*} \mu_Q, \qquad \nu_{R_n} \xrightarrow{*} \mu_R, \tag{2.16}$$

where the convergence is in the sense of weak convergence of measures, i.e.,* $\mu_n \xrightarrow{*} \mu$ *if* $\int f d\mu_n \to \int f d\mu$ *for every bounded continuous function* f*. Furthermore, we have*

$$\nu_{E_n} \to \mu_E, \tag{2.17}$$

in the sense that

$$\lim_{n\to\infty} \int f(s) d\nu_{E_n}(s) = \int f(s) d\mu_E(s)$$

for every continuous function f *such that* $f(s) = \mathcal{O}(s^{-2})$ *as* $s \to \infty$*.*

The convergence of the zero counting measures is due to Stahl [35]. The theorem shows that the measures μ_P, μ_Q, μ_R, and μ_E agree with the measures that Stahl introduced in a different way.

In contrast to the measures ν_{P_n}, ν_{Q_n}, and ν_{R_n} which are probability measures, the measures ν_{E_n} have infinite mass. They also have unbounded support. As a result, the proof of the limit (2.17) is more involved than that of (2.16).

2.3 The g-functions

For the strong asymptotic results we need the log-transforms (or complex logarithmic potentials) of the measures μ_P, μ_Q, and μ_R.

Definition 2.6. We introduce three functions

$$g_P(z) = \int_{\Gamma_P} \log(z-s)\, d\mu_P(s), \qquad z \in \mathbb{C} \setminus \Gamma_P, \tag{2.18}$$

$$g_Q(z) = \int_{\Gamma_Q} \log(z-s)\, d\mu_Q(s), \qquad z \in \mathbb{C} \setminus \Gamma_Q, \tag{2.19}$$

$$g_R(z) = \int_{\Gamma_R} \log(z-s)\, d\mu_R(s), \qquad z \in \mathbb{C} \setminus \Gamma_R, \tag{2.20}$$

which are defined modulo $2\pi i$.

Thus g_P, g_Q, and g_R are multivalued functions, depending on the specific choice of the branches of the logarithmic functions. Our results will involve expressions like e^{ng_P} and e^{ng_R}, and then the multivaluedness will play no role.

2.4 Functions φ_P and φ_R

Two other important functions are the functions φ_P and φ_R given by

$$\varphi_P(z) = \frac{3}{2} \int_{z_1}^{z} (\psi_Q - \psi_P)(s)\, ds, \tag{2.21}$$

and

$$\varphi_R(z) = \frac{3}{2} \int_{z_3}^{z} (\psi_Q - \psi_R)(s)\, ds. \tag{2.22}$$

The paths of integration in (2.21) and (2.22) are in $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R \cup \{0\})$. The functions φ_P and φ_R are multivalued but the real parts are well-defined. From Proposition 2.2 we know that $\operatorname{Re} \varphi_P = 0$ on the curves Γ_P, Γ_P^*, $\Gamma_{E,1}$, and $\Gamma_{E,2}$, and that $\operatorname{Re} \varphi_R = 0$ on Γ_R, Γ_R^*, $\Gamma_{E,1}$, and $\Gamma_{E,2}$. We collect the main properties of φ_P and φ_R in the following lemma.

Lemma 2.7. (a) *The real part of φ_P is zero exactly on Γ_P, Γ_P^*, $\Gamma_{E,1}$, and $\Gamma_{E,2}$.*

The real part of φ_P is negative in $D_{\infty,P} \cup D_P^*$, and it is positive in the remaining part of the plane.

(b) *The real part of φ_R is zero exactly on Γ_R, Γ_R^*, $\Gamma_{E,3}$, and $\Gamma_{E,4}$.*

The real part of φ_R is negative in $D_{\infty,R} \cup D_R^*$, and it is positive in the remaining part of the plane.

(c) *On the imaginary axis, we have $\operatorname{Re} \varphi_P = \operatorname{Re} \varphi_R$. We have $\operatorname{Re} \varphi_P < \operatorname{Re} \varphi_R$ in the left half-plane, and $\operatorname{Re} \varphi_P > \operatorname{Re} \varphi_R$ in the right half-plane.*

Lemma 2.7 is proved in Section 4.5.

2.5 Strong asymptotics away from the zeros

Now we can state the strong asymptotic results for the polynomials P_n, Q_n, R_n, and the remainder function E_n. Recall that the three polynomials are given by (1.4), (1.5) and (1.6) with the constant C as in (1.7) so that Q_n is a monic polynomial. Then P_n and R_n both have leading coefficients $(-\frac{1}{2})^{n+1}$, and we will state our asymptotic results for the monic polynomials $(-2)^{n+1} P_n(z)$ and $(-2)^{n+1} R_n(z)$. Throughout the rest of the paper, we will use the function $\sqrt{3w^4 + 1}$ which branches at the four points w_k given in (2.5). We choose as cuts for this function the two curves $\psi_{P+}(\Gamma_P)$ and $\psi_{R+}(\Gamma_R)$ (see Figure 3), and assume that it is positive for large positive w. So, in particular we have that $\sqrt{3w^4 + 1} = -1$ for $w = 0$.

The following theorem gives the strong asymptotics of the polynomials P_n, Q_n, R_n and the remainder term E_n away from their zeros. These results are due to Stahl [35], but we will give independent proofs below.

Theorem 2.8. *With the functions defined above, we have*

$$(-2)^{n+1} P_n(z) = \frac{2 e^{n g_P(z)}}{\sqrt{3\psi_P^4(z) + 1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \tag{2.23}$$

uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma_P$,

$$(-2)^{n+1} R_n(z) = \frac{2 e^{n g_R(z)}}{\sqrt{3\psi_R^4(z) + 1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \tag{2.24}$$



Figure 6: The shaded region is where $\operatorname{Re}\varphi_P$ is negative. It consists of the two parts $D_{\infty,P}$ and D_P^*, where D_P^* is bounded by Γ_P and Γ_P^*.

uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma_R$, and

$$Q_n(z) = \begin{cases} -\dfrac{e^{ng_Q(z)}}{\sqrt{3\psi_Q^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_\infty \\ \dfrac{e^{ng_Q(z)}}{\sqrt{3\psi_P^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_P \\ \dfrac{e^{ng_Q(z)}}{\sqrt{3\psi_R^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_R, \end{cases} \tag{2.25}$$

uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma_Q$. Furthermore we have

$$E_n(z) = \begin{cases} -\left(-\dfrac{1}{2}\right)^n \dfrac{e^{n(g_R(z)+3z)}}{\sqrt{3\psi_R^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_{\infty,R} \\ -\left(-\dfrac{1}{2}\right)^n \dfrac{e^{n(g_P(z)-3z)}}{\sqrt{3\psi_P^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_{\infty,P} \\ -\dfrac{z^{3n} e^{-n(g_P(z)+g_R(z))}}{\sqrt{3\psi_Q^4(z)+1}} \left(1 + \mathcal{O}\left(\dfrac{1}{n}\right)\right) & \text{for } z \in D_P \cup D_R \cup D_{\infty,U} \cup D_{\infty,L} \\ & \quad \cup\, \Gamma_Q \setminus \{z_1, z_2, z_3, z_4\}, \end{cases} \tag{2.26}$$

uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma_E$.

2.6 Asymptotics near the curves Γ_P, Γ_Q, Γ_R and Γ_E

We will also obtain uniform asymptotics near the curves Γ_P, Γ_Q, Γ_R and Γ_E, as well as in neighborhoods of the branch points z_k. We start with the polynomials Q_n and the remainder function E_n.

Theorem 2.9. *Uniformly for z in compact subsets of the region*

$$(\Gamma_Q \setminus \{z_1, z_2, z_3, z_4\}) \cup D_P \cup D_R \cup D_{\infty,U} \cup D_{\infty,L},$$

see Figure 5, we have

$$\begin{aligned} Q_n(z) \quad &= \quad z^{3n} e^{-n(g_P(z)+g_R(z))} \left[\frac{e^{-2n\varphi_P(z)}}{\sqrt{3\psi_P^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \right. \\ & \left. - \frac{1}{\sqrt{3\psi_Q^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) + \frac{e^{-2n\varphi_R(z)}}{\sqrt{3\psi_R^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \right]. \end{aligned} \tag{2.27}$$

Uniformly for z in compact subsets of D_∞ (which includes the arcs Γ_E but not the branch points z_1, z_2, z_3, z_4), we have

$$\begin{aligned} E_n(z) \quad &= \quad -z^{3n} e^{-n(g_P(z)+g_R(z))} \left[\frac{e^{-2n\varphi_P(z)}}{\sqrt{3\psi_P^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \right. \\ & \left. + \frac{1}{\sqrt{3\psi_Q^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) + \frac{e^{-2n\varphi_R(z)}}{\sqrt{3\psi_R^4(z)+1}} \left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) \right]. \end{aligned} \tag{2.28}$$

Note that the asymptotic formula (2.27) (resp. (2.28)) holds in particular on Γ_Q (resp. Γ_E), away from the branch points z_1, z_2, z_3 and z_4, that is, on the curve where the zeros of Q_n (resp. E_n) accumulate.

It may be checked that the three terms in (2.27) are analytic continuations of the different asymptotic formulas we have in (2.25). For $z \in D_\infty$, we have $g_Q(z) = -g_P(z) - g_R(z) + 3\log z$ by Corollary 4.11, and then (2.27) reduces to the first formula in (2.25) since $\operatorname{Re}\varphi_P(z) > 0$ and $\operatorname{Re}\varphi_R(z) > 0$ in $D_{\infty,U}$ and $D_{\infty,L}$. For $z \in D_P$, we have $g_Q(z) = -g_P(z) - g_R(z) + 3\log z - 2\varphi_P(z)$ and we obtain the second formula in (2.25). Finally, for $z \in D_R$, we have $g_Q(z) = -g_P(z) - g_R(z) + 3\log z - 2\varphi_R(z)$, and we get the third formula in (2.25).

On Γ_Q two of the terms in (2.27) have comparable absolute values. This causes the zeros of Q_n to be close to Γ_Q. At the points $\pm iy^*$, all three terms have comparable absolute values. Similarly, on Γ_E two of the terms in (2.28) have comparable absolute values, which causes the zeros of E_n in D_∞ to be close to Γ_E.

For the polynomials P_n and R_n we have an asymptotic formula on Γ_P and Γ_R, respectively, away from the branch points, which now consists of two terms.

Theorem 2.10. *Uniformly for z in compact subsets of the region*

$$(\Gamma_P \setminus \{z_1, z_2\}) \cup D_{\infty,P} \cup D_P^*$$

(this is the shaded region in Figure 6), we have

$$(-2)^{n+1}P_n(z) = e^{ng_P(z)}\left[\frac{2}{\sqrt{3\psi_P^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \pm \frac{2e^{2n\varphi_P(z)}}{\sqrt{3\psi_Q^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\right] \tag{2.29}$$

where the $+$sign holds in $D_P \cup D_R$ and the $-$sign holds in D_∞.

Uniformly for z in compact subsets of the domain

$$(\Gamma_R \setminus \{z_3, z_4\}) \cup \{z \in \mathbb{C} \mid \operatorname{Re}\varphi_R(z) < 0\}$$

we have

$$(-2)^{n+1}R_n(z) = e^{ng_R(z)}\left[\frac{2}{\sqrt{3\psi_R^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \pm \frac{2e^{2n\varphi_R(z)}}{\sqrt{3\psi_Q^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\right] \tag{2.30}$$

where the $+$sign holds in $D_R \cup D_P$ and the $-$sign holds in D_∞.

For z away from Γ_P so that $\operatorname{Re}\varphi_P(z) < 0$, the asymptotic formula (2.29) reduces to (2.23). On Γ_P we have $\operatorname{Re}\varphi_P(z) = 0$, and then the two terms in (2.29) are of comparable magnitude. For $z \in \Gamma_P$, (2.29) can be re-written as

$$(-2)^{n+1}P_n(z) = \frac{2e^{ng_{P-}(z)}}{\sqrt{3\psi_{P-}^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) + \frac{2e^{ng_{P+}(z)}}{\sqrt{3\psi_{P+}^4(z)+1}}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right).$$

Similar remarks hold for (2.30).

2.7 Asymptotics near the branch points

Near the branch points, the asymptotic formulas involve the Airy function Ai, which is the solution of the differential equation

$$y''(z) = zy(z),$$

satisfying

$$\operatorname{Ai}(z) = \frac{1}{2\sqrt{\pi}} z^{-1/4} e^{-\frac{2}{3}z^{3/2}}\left(1+\mathcal{O}\left(\frac{1}{z^{3/2}}\right)\right)$$

as $z \to \infty$ with $|\arg z| < \pi$, see, e.g., [28] for more details. We only deal with the asymptotic behavior of P_n, Q_n and E_n near z_1. Similar results can be given for the behavior near the other branch points. Here we will take the branch of the function $\varphi_P(z)$ which is 0 at $z = z_1$. So it behaves like

$$\varphi_P(z) = c(z-z_1)^{3/2} + \mathcal{O}\left((z-z_1)^{5/2}\right)$$

as $z \to z_1$, with $c \neq 0$. Then we define the function

$$f_1(z) = \left[\frac{3}{2}\varphi_P(z)\right]^{2/3} \tag{2.31}$$

which is analytic for z in a neighborhood of z_1. We take that 2/3rd power so that $f_1(z)$ is real and negative for $z \in \Gamma_P$. Then

$$f_1(z) = c_1(z - z_1) + \mathcal{O}\left((z - z_1)^2\right) \tag{2.32}$$

as $z \to z_1$ with some constant $c_1 \neq 0$. Explicit calculations show that

$$c_1 = f_1'(z_1) = 2^{1/3}3^{5/12}e^{-\frac{7}{36}\pi i}. \tag{2.33}$$

Theorem 2.11. *There is a $\delta > 0$, such that we have, uniformly for $|z - z_1| < \delta$,*

$$\begin{aligned}(-2)^{n+1}P_n(z) = \sqrt{\pi}e^{(n+1)(g_P(z)+\varphi_P(z))} & \\ \Bigg[n^{1/6}h_1(z)\operatorname{Ai}\left((n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) & \\ +n^{-1/6}h_2(z)\operatorname{Ai}'\left((n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\Bigg], &\end{aligned} \tag{2.34}$$

$$\begin{aligned}(-2)^{n+1}Q_n(z) = \sqrt{\pi}e^{3z}e^{(n+1)(g_P(z)+\varphi_P(z)-3z)} & \\ \Bigg[n^{1/6}h_1(z)e^{-2\pi i/3}\operatorname{Ai}\left(e^{-2\pi i/3}(n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) & \\ +n^{-1/6}h_2(z)e^{2\pi i/3}\operatorname{Ai}'\left(e^{-2\pi i/3}(n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\Bigg], &\end{aligned} \tag{2.35}$$

and

$$\begin{aligned}(-2)^{n+1}E_n(z) = -\sqrt{\pi}e^{3z}e^{(n+1)(g_P(z)+\varphi_P(z)-3z)} & \\ \Bigg[n^{1/6}h_1(z)e^{2\pi i/3}\operatorname{Ai}\left(e^{2\pi i/3}(n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) & \\ +n^{-1/6}h_2(z)e^{-2\pi i/3}\operatorname{Ai}'\left(e^{2\pi i/3}(n+1)^{2/3}f_1(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\Bigg], &\end{aligned} \tag{2.36}$$

where h_1 and h_2 are two analytic functions without zeros in $|z - z_1| < \delta$, which have explicit expressions

$$h_1(z) = \left(N_{21}(z) + iz^{-1}e^{-3z}N_{22}(z)\right) f_1(z)^{1/4}, \tag{2.37}$$

with the branch of the fourth root in $f_1(z)^{1/4}$ taken with a cut along Γ_P, and

$$h_2(z) = \left(-N_{21}(z) + iz^{-1}e^{-3z}N_{22}(z)\right) f_1(z)^{-1/4}. \tag{2.38}$$

Here

$$N_{21}(z) = \frac{2e^{-g_P(z)}}{\sqrt{3\psi_P^4(z)+1}}, \qquad N_{22}(z) = -\frac{e^{g_P(z)+g_R(z)}}{z^2\sqrt{3\psi_Q^4(z)+1}}. \tag{2.39}$$

We use N_{21} and N_{22} to denote these functions, since in what follows they will appear as entries in a matrix N. Note that the function $g_P + \varphi_P$ is analytic near $z = z_1$.

From the asymptotics near the branch points, one can deduce the behavior of the extreme zeros of P_n, Q_n, R_n and E_n near the branch points. We only state the result for the zeros of P_n, Q_n and E_n near z_1. Recall that the Airy function Ai has only negative real zeros, which we denote by $0 > -\iota_1 > -\iota_2 > \cdots > -\iota_\nu > \cdots$.

Corollary 2.12. *Let $z^P_{\nu,n}$, $\nu = 1,\dots,n$, be the zeros of P_n, ordered by increasing distance to z_1. Then for every $\nu \in \mathbb{N}$, we have*

$$\begin{aligned} z^P_{\nu,n} &= z_1 - \frac{\iota_\nu}{f_1'(z_1)} n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \\ &= z_1 + 2^{-1/3} 3^{-5/12} e^{-\frac{29}{36}\pi i} \iota_\nu n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \end{aligned} \tag{2.40}$$

as $n \to \infty$. Let $z^Q_{\nu,n}$, $\nu = 1,\dots,n$, be the zeros of Q_n, ordered by increasing distance to z_1. Then for every $\nu \in \mathbb{N}$,

$$\begin{aligned} z^Q_{\nu,n} &= z_1 - e^{2\pi i/3} \frac{\iota_\nu}{f_1'(z_1)} n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \\ &= z_1 + 2^{-1/3} 3^{-5/12} e^{-\frac{5}{36}\pi i} \iota_\nu n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \end{aligned} \tag{2.41}$$

as $n \to \infty$. Let $z^E_{\nu,n}$, $\nu = 1,\dots,n$, be the zeros of E_n, ordered by increasing distance to z_1. Then for every $\nu \in \mathbb{N}$,

$$\begin{aligned} z^E_{\nu,n} &= z_1 - e^{-2\pi i/3} \frac{\iota_\nu}{f_1'(z_1)} n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \\ &= z_1 + 2^{-1/3} 3^{-5/12} e^{\frac{19}{36}\pi i} \iota_\nu n^{-2/3} + \mathcal{O}\left(\frac{1}{n}\right), \end{aligned} \tag{2.42}$$

as $n \to \infty$.

2.8 Convergence of the Hermite–Padé approximants

The polynomials P_n, Q_n, R_n satisfy

$$P_n(z) + Q_n(z) e^{3nz} + R_n(z) e^{6nz} = \mathcal{O}\left(z^{3n+2}\right),$$

as $z \to 0$. Replacing the $\mathcal{O}$ term by 0, one might expect that the quadratic equation

$$P_n(z) + Q_n(z) X_n(z) + R_n(z) X_n^2(z) = 0 \tag{2.43}$$

has a solution

$$X_n(z) = \frac{-Q_n(z) \pm \sqrt{Q_n^2(z) - 4P_n(z)R_n(z)}}{2R_n(z)} \tag{2.44}$$

which is close to the exponential e^{3nz}. This is indeed true for z in the domains D_P and D_R.

Theorem 2.13. *For $z \in D_P \cup D_R$, there is a solution to (2.43) such that*

$$X_n(z) = e^{3nz}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right). \tag{2.45}$$

The $\mathcal{O}$-term is uniform on compact subsets of $D_P \cup D_R$.

Remark 2.14. The domain of convergence does *not* include the imaginary axis. On the imaginary axis one has roots of $Q_n^2 - 4P_nR_n$ which are branch points of X_n, accumulating along the interval $[-iy^*, iy^*]$. Hence, convergence does not take place in a neighborhood of 0 even though the approximants come from a Hermite–Padé expansion around 0.

Remark 2.15. It turns out that Q_n^2 dominates P_nR_n in the region $D_P \cup D_R$. If we agree to take the square root in (2.44) so that

$$\sqrt{Q_n^2(z) - 4P_n(z)R_n(z)} \simeq Q_n(z),$$

then in (2.44) we have to take the $+$ sign in D_P and the $-$ sign in D_R. So

$$\begin{aligned} X_n(z) &= \frac{-Q_n(z) + \sqrt{Q_n^2(z) - 4P_n(z)R_n(z)}}{2R_n(z)} \\ &= \frac{-2P_n(z)}{Q_n(z) + \sqrt{Q_n^2(z) - 4P_n(z)R_n(z)}} \simeq -\frac{P_n(z)}{Q_n(z)} \qquad \text{for } z \in D_P, \end{aligned}$$

and

$$X_n(z) = \frac{-Q_n(z) - \sqrt{Q_n^2(z) - 4P_n(z)R_n(z)}}{2R_n(z)} \simeq -\frac{Q_n(z)}{R_n(z)} \qquad \text{for } z \in D_R.$$

Remark 2.16. It is possible to show that in D_∞, there is no relative convergence to e^{3nz} of any of the two roots $X_n(z)$. Absolute convergence holds in D_P and in a subdomain of D_R bounded by the curve $\operatorname{Re}(3z + 2\varphi_R(z)) = 0$.

3 Geometry of the problem

In this section we will prove Propositions 2.1 and 2.2 together.

3.1 Proof of Propositions 2.1 and 2.2

We recall that ψ_P, ψ_Q, and ψ_R will be the three inverse mappings of

$$z = z(w) = \frac{w^2 - \frac{1}{3}}{w(w^2-1)}.$$

It is easy to see that $z(w)$ is real if w is real, and that $z(w)$ is purely imaginary if $w \in i\mathbb{R}$. In fact, for $z \in i\mathbb{R}$, there is one solution of the cubic equation (2.4) on the imaginary axis, one in the left half-plane, and one in the right half-plane. We denote these by $\psi_Q(z)$, $\psi_P(z)$, and $\psi_R(z)$, respectively. Then part (b) of Proposition 2.1 holds. We also note that $\psi_Q(0) = \infty$.

These functions have analytic continuations, first to a vertical strip $-\varepsilon < \operatorname{Re} z < \varepsilon$. around the imaginary axis. Then ψ_Q has a pole at 0 and from (2.3) it follows that

$$\psi_Q(z) = \frac{1}{z} + \mathcal{O}(z) \qquad \text{as } z \to 0. \tag{3.1}$$

Near infinity, we have from analyzing (2.3)

$$\psi_P(z) = -1 + \frac{1}{3z} + \mathcal{O}\left(\frac{1}{z^2}\right), \tag{3.2}$$

$$\psi_Q(z) = \frac{1}{3z} + \mathcal{O}\left(\frac{1}{z^2}\right), \tag{3.3}$$

$$\psi_R(z) = 1 + \frac{1}{3z} + \mathcal{O}\left(\frac{1}{z^2}\right), \tag{3.4}$$

as $z \to \infty$.

For $0 < \operatorname{Re} z < \varepsilon$, we then have that $\operatorname{Re} \psi_Q(z) > 0$, $\operatorname{Re} \psi_R(z) > 0$, while $\operatorname{Re} \psi_P(z) < 0$. Since for $z \notin i\mathbb{R}$, there can be no solution of (2.4) on the imaginary axis, it then follows by continuity that for every z in the right half-plane, there are two solutions of the cubic equation (2.4) with positive real part, and one solution with negative real part. For z with $\operatorname{Re} z > 0$, we then define $\psi_P(z)$ as the unique solution of (2.4) with negative real part. This defines ψ_P as an analytic function for $\operatorname{Re} z > -\varepsilon$. Similarly, for $\operatorname{Re} z < 0$, we define $\psi_R(z)$ as the unique solution of (2.4) with positive real part, and we have ψ_R as an analytic function for $\operatorname{Re} z < \varepsilon$.

For the moment, we let Γ_P be an arbitrary simple curve that connects the branch points z_1 and z_2, and lies entirely in the left half-plane. We let Γ_R be the mirror image of Γ_P with respect to the imaginary axis and we assume that these cuts determine the sheet structure of the Riemann surface $\mathcal{R}$. Then we can extend ψ_P, ψ_Q, and ψ_R to analytic functions on $\mathbb{C} \setminus \Gamma_P$, $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R)$, and $\mathbb{C} \setminus \Gamma_R$, respectively. So we have part (a) of Proposition 2.1.

Next we want to study curves so that $\frac{3}{2\pi i} \int_{z_1}^{z} (\psi_Q - \psi_P)(s) ds$ is real. If $z = z(t)$ is arclength parametrization of such a curve, then

$$\operatorname{Re} \left[\int_{z_1}^{z(t)} (\psi_Q - \psi_P)(s) ds \right] = 0.$$

which upon differentiating leads to Re $[z'(t)(\psi_Q - \psi_P)(z(t))] = 0$. Since $z'(t) \neq 0$, and $(\psi_Q - \psi_P)(z) \neq 0$ (except at the branch points z_1 and z_2), we find that $-z'(t)^2(\psi_Q - \psi_P)(z(t))^2 > 0$. Thus the curve is what is known in geometric function theory as a trajectory of the quadratic differential

$$-(\psi_Q - \psi_P)^2(z) dz^2, \tag{3.5}$$

see [29, 37]. Now, $(\psi_Q - \psi_P)^2(z)$ is well-defined in the left half-plane, irrespective of the exact choice for Γ_P. It is analytic with simple zeros at the branch points z_1 and z_2, and a double pole at 0.

Trajectories of the quadratic differential (3.5) which start from or end at z_1 or z_2 are called critical trajectories. From the local structure of trajectories of quadratic differentials, it is known that three trajectories emanate from any simple zero. So three critical trajectories emanate from z_1, and we would like to label them by Γ_P, Γ_P^*, and $\Gamma_{E,1}$. The labeling will depend on the global behavior of these trajectories. We consider the global behavior in the second quadrant, which we call G.

It is known that any trajectory in G must begin and end either at z_1, or at infinity, or on the boundary of G. It is also known that there can be no closed Jordan curve consisting of trajectories, since there are no poles of the quadratic differential in G. For these properties of trajectories of quadratic differentials, see [29, Chapter 8] and also [3]. So the three critical trajectories that start at z_1, cannot end at z_1, and they should end in G in one of the following ways

(1) on the negative real axis,

(2) on the positive imaginary axis, or

(3) at infinity.

If a critical trajectory meets the real axis, then by symmetry its mirror image with respect to the real axis will be its continuation to z_2. If there were two critical trajectories that come to the negative real axis, then they both would continue to z_2. Then there would be a closed Jordan curve consisting of trajectories, which would not enclose a pole. This is impossible. Hence there can be at most one critical trajectory that meets the negative real axis.

On the critical trajectories we have that $\varphi_P(z)$, see (2.21), is purely imaginary. Its derivative is $\varphi_P'(z) = \frac{3}{2\pi i}(\psi_Q - \psi_P)(z)$, and

$$\operatorname{Re}\varphi_P'(z) = \frac{3}{2\pi}\operatorname{Im}(\psi_Q - \psi_P)(z). \tag{3.6}$$

We claim that $\operatorname{Im}\psi_Q(z) = \operatorname{Im}\psi_P(z)$ cannot happen for z on the positive imaginary axis. To prove this claim, let us assume that for $z = iy$ with $y > 0$, we have $\operatorname{Im}\psi_Q(z) = \operatorname{Im}\psi_P(z)$. Note that ψ_Q maps the positive imaginary axis to the negative imaginary axis. So we have $\psi_Q(iy) = -iv$ with $v > 0$ and

$$y = \frac{v^2 + \frac{1}{3}}{v(v^2+1)}. \tag{3.7}$$

Recall that $\psi_P(z)$ is always in the left half-plane. So we have $\psi_P(iy) = -u - iv$ with $u > 0$, and by symmetry, we then have $\psi_R(iy) = u - iv$. Since ψ_P, ψ_Q, and ψ_R are the three solutions of (2.4), we have

$$\psi_P(z) + \psi_Q(z) + \psi_R(z) = \frac{1}{z}, \qquad z \in \mathbb{C}.$$

Then for $z = iy$, this leads to $-3iv = \frac{1}{iy}$, so that $y = \frac{1}{3v}$. However, this is incompatible with (3.7) if $v \neq 0$. This contradiction proves the claim that $\operatorname{Im}\psi_Q(z) = \operatorname{Im}\psi_P(z)$ does not happen for z on the positive imaginary axis. As a result we see from (3.6) that the real part of $\varphi_P(z)$ is strictly monotone as z varies over the positive imaginary axis. Thus there can be at most one value $y^* > 0$ so that $\operatorname{Re}\varphi_P(iy^*) = 0$. Since $\operatorname{Re}\varphi_P = 0$ on the critical trajectories, we see that there can be at most one critical trajectory that meets the positive imaginary axis.

Now consider trajectories that end at infinity. At infinity, we have $-(\psi_Q - \psi_P)^2(\infty) = -1$. This means that the quadratic differential has a pole of order 4 at infinity, see [37], and all trajectories that extend to infinity arrive there with a vertical tangent. Assume that two critical trajectories extend to infinity in G. Then these trajectories are the boundary of a region G' in G. Any trajectory in G' begins and ends at infinity with a vertical tangent. However, according to [37, Theorem 7.4] a pole of order 4 has a neighborhood, so that any closed trajectory lying entirely in that neighborhood begins and ends at the pole, but from opposite directions. This is a contradiction, since there will be trajectories in G' that are arbitrarily close to infinity. This contradiction shows that there can be at most one critical trajectory that extends to infinity.

So now we proved that each of the possible ways (1), (2), and (3) that a critical trajectory can end in G, happens at most once. Since there are three critical trajectories, it follows that each possibility happens exactly once. So there is a unique critical trajectory that meets the negative real axis. This trajectory continues to z_2, and we call it Γ_P. We then have that part (c) of Proposition 2.1 is satisfied.

The critical trajectory that meets the positive imaginary axis is called Γ_P^*, and the critical trajectory that extends to infinity is called $\Gamma_{E,1}$. By symmetry, we also find the trajectory $\Gamma_{E,2}$ that emanates from z_2 and extends to infinity.

By symmetry with respect to the imaginary axis, we also find the contours Γ_R, $\Gamma_{E,3}$, $\Gamma_{E,4}$, and Γ_R^* in the right half-plane, so that for z on these contours we have that $\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q -$

$\psi_R)(s)ds$ is real. This establishes part (d) of Proposition 2.1 and this proposition is now proved completely.

Large parts of Proposition 2.2 have also been proved. All statements regarding Γ_P, Γ_R, $\Gamma_{E,k}$, $k = 1, 2, 3, 4$ are proved. We also know that Γ_P^* and Γ_R^* meet at the point iy^* on the positive imaginary axis. What remains is to investigate how Γ_P^* and Γ_R^* continue beyond iy^*. The following lemma will be useful.

Lemma 3.1. *The real part of* φ_P *is strictly positive on* Γ_R*,* $\Gamma_{E,3}$*, and* $\Gamma_{E,4}$*. The real part of* φ_R *is strictly positive on* Γ_P*,* $\Gamma_{E,1}$*, and* $\Gamma_{E,2}$*.*

We delay the proof of this lemma to Section 4.4, since its proof is easier once we have the functions g_P and g_R and know some of their properties. The reader may verify that the proofs that follow in Section 4 do not depend on the curves Γ_P^* and Γ_R^*.

Assuming the lemma we can complete the proof of Proposition 2.2 as follows. Consider the quadratic differential $-(\psi_Q - \psi_P)(z)^2 dz^2$ in the domain G_1 bounded by $\Gamma_{E,4}$, the curve Γ_R in the upper half-plane and the real interval $(-\infty, x^*]$ where x^* is the point where Γ_R intersects the real axis. The critical trajectory Γ_P^* has to end in G_1 in one of the following ways

(1) on the real axis,

(2) on Γ_R or $\Gamma_{E,4}$,

(3) at infinity.

It cannot end on Γ_R or $\Gamma_{E,4}$ because of Lemma 3.1, and it cannot extend to infinity for the same reason that it could not extend to infinity in the left half-plane. So it has to end on the real axis, but it cannot end on the negative real axis, again for the same reason as before. So it has to end on the positive real axis, somewhere in the open interval $(0, x^*)$. Then by symmetry Γ_P^* continues in the lower half-plane to the branch point z_2 and Γ_P^* does not intersect with Γ_R. This proves part (c) of Proposition 2.2.

By symmetry, also part (d) holds, and this completes the proof of Proposition 2.2.

4 Measures and functions associated with the Riemann surface

In this section we study the measures μ_P, μ_Q, μ_R, and μ_E, and the functions φ_P, φ_R, g_P, g_Q, and g_R. These measures and functions that are associated with the Riemann surface satisfy many relations that will be used in the transformations of the Riemann–Hilbert problem that follow in later sections. We also prove Theorem 2.4 and Lemmas 2.7 and 3.1 in this section.

4.1 Properties of the measures μ_P, μ_R, and μ_E

We start with a lemma.

Lemma 4.1. *We have*

$$\frac{3}{2\pi i} \int_{\Gamma_P} (\psi_Q - \psi_P)_+(s)\, ds = 1, \tag{4.1}$$

and

$$\frac{3}{2\pi i} \int_{\Gamma_R} (\psi_Q - \psi_R)_+(s)\, ds = 1. \tag{4.2}$$

Proof. Let γ be a closed contour on the sheet $\mathcal{R}_P$ going around Γ_P once in the positive direction. Then the residue theorem for the exterior of γ gives

$$\frac{3}{2\pi i}\int_\gamma \psi_P(s)\,ds = 1,$$

because ψ_P is analytic outside γ and we have (3.2). If we shrink γ to Γ_P, then the integral becomes

$$\frac{3}{2\pi i}\int_{\Gamma_P} ((\psi_P)_-(s) - (\psi_P)_+(s))\,ds = 1.$$

Taking into account that $(\psi_P)_- = (\psi_Q)_+$, we obtain (4.1). The reasoning is similar for the proof of (4.2), where we use a closed contour going around Γ_R on $\mathcal{R}_R$ and the behavior (3.4) of ψ_R at infinity. ☐

Now we can prove the parts of Theorem 2.4 dealing with μ_P, μ_R and μ_E. We give the proof of the part dealing with μ_Q only after we have a better understanding of the curve Γ_P^* and Γ_R^*, see Proposition 4.8.

Proposition 4.2. *μ_P is a probability measure on Γ_P, μ_R is a probability measure on Γ_R and μ_E is a positive measure on Γ_E.*

Proof. The curve Γ_P is such that for $z \in \Gamma_P$, the integral $\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)_+(s)\,ds$ is real, see Proposition 2.1. For $z = z_1$, it has the value 0, and for $z = z_2$ it has the value 1 by (4.1). Let $z = z(t)$, $t \in [0, T]$, be arclength parametrization of Γ_P. The derivative of

$$t \mapsto \frac{3}{2\pi i}\int_{z_1}^{z(t)} (\psi_Q - \psi_P)_+(s)\,ds \tag{4.3}$$

is equal to $\frac{3}{2\pi i}\left(\psi_Q(z(t)) - \psi_P(z(t))\right) z'(t)$ and this is different from 0 for $t \in (0, T)$. Thus (4.3) is strictly increasing from 0 for $t = 0$ to 1 for $t = T$. This immediately implies that μ_P defined by (2.12) is a probability measure on Γ_P. Similarly μ_R defined in (2.13) is a probability measure on Γ_R.

For μ_E we observe that (2.15) defines a real measure on Γ_E, since, by Proposition 2.1, $\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)(s)ds$ is real for $z \in \Gamma_{E,1} \cup \Gamma_{E,2}$, and $\frac{3}{2\pi i}\int_{z_3}^{z}(\psi_Q - \psi_R)(s)ds$ is real for $z \in \Gamma_{E,3} \cup \Gamma_{E,4}$. Using an argument based on arclength parametrization, similar to the one above, we find that on each part $\Gamma_{E,j}$, $j = 1, 2, 3, 4$, the measure μ_E is either positive or negative. Since $\psi_Q(s) - \psi_P(s) = 1 + \mathcal{O}(1/s^2)$ as $s \to \infty$ (see (3.2)–(3.3)), we have for $z \in \Gamma_{E,1}$,

$$\frac{3}{2\pi i}\int_{z_1}^{z}(\psi_Q - \psi_P)(s)\,ds = \frac{3}{2\pi i}(z - z_1) + \mathcal{O}(1) \qquad \text{as } z \to \infty,\ z \in \Gamma_{E,1}.$$

Since $\operatorname{Im}(z - z_1) \to +\infty$ as $z \to \infty$ along $\Gamma_{E,1}$, the integral is positive as $z \to \infty$ along $\Gamma_{E,1}$. As the measure μ_E is of constant sign on $\Gamma_{E,1}$, we may thus deduce that it is positive everywhere on $\Gamma_{E,1}$. The reasoning for $\Gamma_{E,2}$, $\Gamma_{E,3}$, and $\Gamma_{E,4}$ is similar.

This completes the proof of Proposition 4.2. ☐

4.2 Properties of g_P and g_R

The functions g_P and g_R were defined in (2.18) and (2.20). These are multi-valued functions, depending on the choice of the branch of the logarithm $\log(z-s)$, which we assume depends on $s \in \Gamma$ in a continuous way. Since μ_P and μ_R are probability measures, the g-functions are defined modulo $2\pi i$.

Lemma 4.3. *For the derivatives of the functions g_P and g_R we have*

$$g_P'(z) = 3\psi_P(z) + 3, \qquad z \in \mathbb{C} \setminus \Gamma_P, \tag{4.4}$$

and

$$g_R'(z) = 3\psi_R(z) - 3, \qquad z \in \mathbb{C} \setminus \Gamma_R. \tag{4.5}$$

Proof. The derivative of g_P is easily obtained as

$$g_P'(z) = \frac{3}{2\pi i} \int_{\Gamma_P} \frac{1}{z-s} (\psi_Q - \psi_P)_+(s)\, ds.$$

If γ is a closed contour going around Γ_P on $\mathcal{R}_P$ in the positive direction but with z outside γ, then (since $(\psi_Q)_+ = (\psi_P)_-$)

$$g_P'(z) = \frac{3}{2\pi i} \oint_{\gamma} \frac{\psi_P(s)}{z-s}\, ds.$$

The integral over γ can be calculated with the residue theorem for the exterior of γ, for which there is a residue at z and at ∞ given by (3.2). This proves (4.4). The proof of (4.5) is similar. $\square$

Corollary 4.4. *For $z \in \mathbb{C} \setminus (\Gamma_P \cup \Gamma_R)$, we have*

$$2g_P'(z) + g_R'(z) = \frac{3}{z} + 3 - 3[\psi_Q(z) - \psi_P(z)], \tag{4.6}$$

and

$$2g_R'(z) + g_P'(z) = \frac{3}{z} - 3 - 3[\psi_Q(z) - \psi_R(z)]. \tag{4.7}$$

Proof. We recall that $\psi_P(z)$, $\psi_Q(z)$, and $\psi_R(z)$ are roots of the cubic equation (2.4), so that

$$\psi_P(z) + \psi_Q(z) + \psi_R(z) = \frac{1}{z}. \tag{4.8}$$

Hence by (4.4) and (4.5),

$$\begin{aligned} 2g_P'(z) + g_R'(z) &= 6\psi_P(z) + 6 + 3\psi_R(z) - 3 \\ &= 3[\psi_P(z) + \psi_Q(z) + \psi_R(z)] - 3[\psi_Q(z) - \psi_P(z)] + 3 \\ &= \frac{3}{z} + 3 - 3[\psi_Q(z) - \psi_P(z)]. \end{aligned}$$

A similar computation gives (4.7). $\square$

A useful explicit expression of g_P and g_R in terms of the mapping functions ψ_P and ψ_R is given in the next lemma.

Lemma 4.5. *The functions g_P and g_R have the following explicit expressions in terms of the mapping functions ψ_P and ψ_R,*

$$g_P(z) = 3z(\psi_P(z)+1) - \log[\psi_P(z)(\psi_P^2(z)-1)] - 1 + \log\left(\frac{2}{3}\right), \text{ for } z \in \mathbb{C}\setminus\Gamma_P, \tag{4.9}$$

and

$$g_R(z) = 3z(\psi_R(z)-1) - \log[\psi_R(z)(\psi_R^2(z)-1)] - 1 + \log\left(\frac{2}{3}\right), \text{ for } z \in \mathbb{C}\setminus\Gamma_R. \tag{4.10}$$

Proof. We let $z \in \mathbb{C}\setminus\Gamma_P$ and put $w = \psi_P(z)$. Taking a derivative of (2.3), we find, since $w'(z) = \frac{1}{z'(w)}$,

$$3 = -\left(\frac{1}{w^2} + \frac{1}{(w-1)^2} + \frac{1}{(w+1)^2}\right) w'.$$

Thus

$$\begin{aligned} 3\psi_P(z) + 3 &= -(w+1)\left(\frac{1}{w^2} + \frac{1}{(w-1)^2} + \frac{1}{(w+1)^2}\right) w' \\ &= -\left(\frac{1}{w} + \frac{1}{w^2} + \frac{1}{w-1} + \frac{2}{(w-1)^2} + \frac{1}{w+1}\right) w' \\ &= \frac{d}{dz}\left[-\log w(w^2-1) + \frac{1}{w} + \frac{2}{w-1}\right]. \end{aligned}$$

By (4.4), we then see that

$$g_P(z) = -\log w(w^2-1) + \frac{1}{w} + \frac{2}{w-1} + C$$

for some constant C. The constant can be determined from the behavior for $z \to \infty$, since $g_P(z) = \log z + \mathcal{O}(1/z)$, and $w = \psi_P(z) = -1 + \frac{1}{3z} + \mathcal{O}(1/z^2)$. The result is that $C = 2 + \log\frac{2}{3}$. Using (2.3) we then find (4.9).

The proof of (4.10) follows along similar lines. ☐

4.3 Jump properties of g_P and g_R

We recall that the functions φ_P and φ_R were introduced in (2.21) and (2.22). The next lemma connects these functions with g_P and g_R. It will be frequently used in what follows. Throughout the rest of the paper we use ℓ to denote the constant

$$\ell = \log 2 - \pi i. \tag{4.11}$$

Lemma 4.6. *For $z \in \mathbb{C}\setminus(\Gamma_P \cup \Gamma_R)$, we have*

$$2g_P(z) + g_R(z) = 3\log z + 3z - 2\varphi_P(z) + \ell, \tag{4.12}$$

$$2g_R(z) + g_P(z) = 3\log z - 3z - 2\varphi_R(z) + \ell. \tag{4.13}$$

On the contours we have

$$g_{P+}(z) + g_{P-}(z) + g_R(z) = 3\log z + 3z + \ell, \qquad z \in \Gamma_P, \tag{4.14}$$

$$g_P(z) + g_{R+}(z) + g_{R-}(z) = 3\log z - 3z + \ell, \qquad z \in \Gamma_R, \tag{4.15}$$

and

$$g_{P+}(z) - g_{P-}(z) = -\varphi_{P+}(z) + \varphi_{P-}(z) = -2\varphi_{P+}(z) = 2\varphi_{P-}(z), \qquad z \in \Gamma_P, \tag{4.16}$$

$$g_{R+}(z) - g_{R-}(z) = -\varphi_{R+}(z) + \varphi_{R-}(z) = -2\varphi_{R+}(z) = 2\varphi_{R-}(z), \qquad z \in \Gamma_R. \tag{4.17}$$

Proof. Integrating (4.6) from z_1 to z over some path in $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R \cup \{0\})$, we get

$$2g_P(z) + g_R(z) - 2g_P(z_1) - g_R(z_1) = 3\log z - 3\log z_1 + 3(z - z_1) - 2\varphi_P(z),$$

so that (4.12) holds with constant

$$\ell = 2g_P(z_1) + g_R(z_1) - 3\log z_1 - 3z_1.$$

Using the explicit expressions (4.9) and (4.10) for g_P and g_R we are able to show that ℓ is equal to (4.11). To obtain (4.13) we integrate (4.7) from z_3 to z. It turns out that the constant of integration is again given by (4.11).

Next, we use Lemma 4.3 to find

$$g_{P+}(z) + g_{P-}(z) + g_R(z) = 3\int_{z_1}^{z} (\psi_{P+} + \psi_{P-} + \psi_R)(s)\, ds + 3(z - z_1), \qquad z \in \Gamma_P.$$

On Γ_P we have $\psi_{P-}(s) = \psi_{Q+}(s)$, so that

$$g_{P+}(z) + g_{P-}(z) + g_R(z) = 3\int_{z_1}^{z} (\psi_P + \psi_Q + \psi_R)_+(s)\, ds + 3z - 3z_1, \qquad z \in \Gamma_P.$$

Since $\psi_P(s) + \psi_Q(s) + \psi_R(s) = \frac{1}{s}$, we obtain

$$g_{P+}(z) + g_{P-}(z) + g_R(z) = 3\log z - 3\log z_1 + 3z - 3z_1 = 3\log z - 3z + \ell, \qquad z \in \Gamma_P.$$

This proves (4.14). A similar computation leads to (4.15).

Finally, if we take (4.12) on the + and −sides of Γ_P and subtract, we get

$$2g_{P+}(z) - 2g_{P-}(z) = -2\varphi_{P+}(z) + 2\varphi_{P-}(z), \qquad z \in \Gamma_P.$$

This gives (4.16), since $\varphi_{P+}(z) = -\varphi_{P-}(z)$. Similarly we find (4.17). $\square$

Since $g_P(z) = \log z + \mathcal{O}(1/z)$ and $g_R(z) = \log z + \mathcal{O}(1/z)$ as $z \to \infty$, we get from (4.12)–(4.13) that

$$2\varphi_P(z) = 3z + \ell + \mathcal{O}\left(\frac{1}{z}\right) \qquad \text{as } z \to \infty, \tag{4.18}$$

and

$$2\varphi_R(z) = -3z + \ell + \mathcal{O}\left(\frac{1}{z}\right) \qquad \text{as } z \to \infty. \tag{4.19}$$

We also see from (4.12) and (4.13) that φ_P and φ_R are multivalued functions, which are defined modulo πi, since g_P and g_R are defined modulo $2\pi i$.

4.4 Proof of Lemma 3.1

Now we are in a position to prove Lemma 3.1 which was needed in Section 3 to complete the discussion on the contours Γ_P^* and Γ_R^*.

Proof. Let γ be the contour which consists of Γ_R, $\Gamma_{E,3}$ and $\Gamma_{E,4}$. We define a function G_R on the right half-plane $\{\operatorname{Re} z > 0\}$ by

$$G_R(z) = \begin{cases} g_R(z) & \text{to the left of } \gamma, \\ g_R(z) + 2\varphi_R(z) & \text{to the right of } \gamma. \end{cases}$$

This function is analytic across Γ_R because of the jump relation (4.17) of g_R. The function $u(z) = \operatorname{Re} G_R(z)$ is therefore harmonic in $\{\operatorname{Re} z > 0\} \setminus (\Gamma_{E,3} \cup \Gamma_{E,4})$. On $\Gamma_{E,3}$ and $\Gamma_{E,4}$ we have $\operatorname{Re} \varphi_R = 0$, and thus u is continuous in the full right half-plane. Furthermore $\operatorname{Re} \varphi_R < 0$ on the right of γ. Since g_R is analytic across $\Gamma_{E,3}$ and $\Gamma_{E,4}$ it then easily follows that u is superharmonic in $\{\operatorname{Re} z > 0\}$. Near infinity we have

$$u(z) = \begin{cases} \operatorname{Re} \log z + \mathcal{O}(1/z) & \text{as } z \to \infty \text{ to the left of } \gamma, \\ \operatorname{Re} (-3z + \log z + \ell) + \mathcal{O}(1/z) & \text{as } z \to \infty \text{ to the right of } \gamma. \end{cases}$$

By symmetry $\operatorname{Re} g_P(z) = \operatorname{Re} g_R(z)$ on the imaginary axis. Thus $\operatorname{Re} g_P(z) - 3\operatorname{Re} z$ is harmonic in the right half plane, $\operatorname{Re} g_P(z) - 3\operatorname{Re} z = u(z)$ on the imaginary axis, and $u(z) - \operatorname{Re} g_P(z) + 3\operatorname{Re} z$ is bounded from below as $z \to \infty$. Then it follows from the minimum principle for superharmonic functions that $u(z) - \operatorname{Re} g_P(z) + 3\operatorname{Re} z \geq 0$ in the right half-plane. In particular this gives $\operatorname{Re} (g_R(z) - g_P(z) + 3z) \geq 0$ to the left of γ and $\operatorname{Re} (g_R(z) - g_P(z) + 2\varphi_R(z)) \geq 0$ to the right of γ. Taking the difference between (4.13) and (4.12) gives $g_R(z) - g_P(z) = -6z - 2\varphi_R(z) + 2\varphi_P(z)$. Then we see that $\operatorname{Re} (-3z - 2\varphi_R(z) + 2\varphi_P(z)) \geq 0$ to the left of γ and $\operatorname{Re} (-3z + 2\varphi_P(z)) \geq 0$ to the right of γ. Letting now $z \to \gamma$ from either side, we get, since $\operatorname{Re} \varphi_R(z) = 0$ on γ, that $2\operatorname{Re} \varphi_P(z) \geq 3\operatorname{Re} z > 0$ on Γ_R, $\Gamma_{E,3}$, and $\Gamma_{E,4}$.

The proof that $\operatorname{Re} \varphi_R > 0$ on Γ_P, $\Gamma_{E,1}$, and $\Gamma_{E,2}$ is similar, and also follows because of symmetry. □

4.5 Proof of Lemma 2.7

Proof. By Proposition 2.1 and the definition of φ_P, we have that $\operatorname{Re} \varphi_P = 0$ on Γ_P, Γ_P^*, $\Gamma_{E,1}$, and $\Gamma_{E,2}$.

We know that $\operatorname{Re} \varphi_P$ is a harmonic function in $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R \cup \{0\})$. Since $\psi_Q(z) \sim 1/z$ as $z \to 0$, it easily follows from (2.21) that $\operatorname{Re} \varphi_P(z) \to -\infty$ as $z \to 0$. Then by the maximum principle for harmonic functions we get that $\operatorname{Re} \varphi_P < 0$ on D_P^*.

As $z \to \infty$, we have (4.18). On the unbounded curves $\Gamma_{E,1}$ and $\Gamma_{E,2}$ we have $\operatorname{Re} \varphi_P = 0$. From (4.18) it then follows that these curves are asymptotic to the vertical line $\operatorname{Re} (3z + \ell) = 0$, which is $\operatorname{Re} z = -\frac{1}{3} \log 2$. As $z \to \infty$ in the unbounded domain $D_{\infty,P}$ we then have $\limsup \operatorname{Re} (3z + \ell) \leq 0$, so that $\limsup \operatorname{Re} \varphi_P(z) \leq 0$ by (4.18). Again it follows by the maximum principle for harmonic functions that $\operatorname{Re} \varphi_P < 0$ on $D_{\infty,P}$.

For the remaining domain $G = \mathbb{C} \setminus (\overline{D_P^* \cup D_{\infty,P}})$, we have that $\operatorname{Re} \varphi_P$ is harmonic on $G \setminus \Gamma_R$, with $\liminf \varphi_P(z) \geq 0$ as $z \to \infty$ with $z \in G$ by (4.18), and $\operatorname{Re} \varphi_P > 0$ on Γ_R by Lemma 3.1. Thus again by the maximum principle, $\operatorname{Re} \varphi_P > 0$ on G. This completes the proof of part (a) of Lemma 2.7.

The proof of part (b) is similar and also follows because of symmetry.

For part (c) we note that $\operatorname{Re}\varphi_P = \operatorname{Re}\varphi_R$ on the imaginary axis because of symmetry. As $z \to \infty$, we have by (4.18) and (4.19) that $\liminf \operatorname{Re}(\varphi_R(z) - \varphi_P(z)) \geq 0$ as $z \to \infty$ in the left half-plane. Since $\operatorname{Re}\varphi_R$ and $\operatorname{Re}\varphi_P$ are harmonic in $\{\operatorname{Re} z < 0\} \setminus \Gamma_P$ and $\operatorname{Re}\varphi_R > \operatorname{Re}\varphi_P$ on Γ_P, we find that $\operatorname{Re}\varphi_P < \operatorname{Re}\varphi_R$ in the left half-plane. Similarly, we have $\operatorname{Re}\varphi_P > \operatorname{Re}\varphi_R$ in the right half-plane.

This completes the proof of Lemma 2.7. ☐

4.6 The measure μ_Q

Now that we have the full curves Γ_P^* and Γ_R^*, we can prove the part of Theorem 2.4 that deals with μ_Q. We start with the analogue of Lemma 4.1.

Lemma 4.7. *We have*

$$\frac{3}{2\pi i}\int_{\Gamma_P^*} (\psi_Q - \psi_P)(s)\,ds = 2, \tag{4.20}$$

and

$$\frac{3}{2\pi i}\int_{\Gamma_R^*} (\psi_Q - \psi_R)(s)\,ds = 2. \tag{4.21}$$

Proof. If we deform the integral (4.20) over Γ_P^* to an integral over Γ_P, then we pick up a residue contribution at 0 due to the pole of ψ_Q at the origin. So by (3.1) and (4.1) we have

$$\frac{3}{2\pi i}\int_{\Gamma_P^*} (\psi_Q - \psi_P)(s)\,ds = 3 - \frac{3}{2\pi i}\int_{\Gamma_P} (\psi_Q - \psi_P)_+(s)\,ds = 2.$$

This gives (4.20) and in a similar way we obtain (4.21). ☐

Now we can prove that μ_Q is a probability measure.

Proposition 4.8. *μ_Q is a probability measure on Γ_Q.*

Proof. We prove first that (2.14) defines a positive measure. On Γ_P^* we have that $\frac{3}{2\pi i}\int_{z_2}^{z}(\psi_Q - \psi_P)(s)ds$ takes on real values, and has the value 0 at $z = z_2$ and the value 2 for $z = z_1$ by (4.20). Since its derivative does not vanish between z_2 and z_1, we find that $\frac{3}{2\pi i}(\psi_Q - \psi_P)(s)ds$ is a positive measure on Γ_P^* of total mass 2. Its restriction to $\Gamma_P^* \cap \{\operatorname{Re} z < 0\}$ is part of the measure μ_Q, and so this part is a positive measure. Similarly, $\frac{3}{2\pi i}(\psi_R - \psi_Q)(s)ds$ is a positive measure on Γ_R^* (due to (4.21)), and so also on $\Gamma_R^* \cap \{\operatorname{Re} z > 0\}$. On the imaginary axis, we have, because of symmetry, that $\psi_P(s) = -\overline{\psi_R(s)}$, which means that $\psi_R(s) - \psi_P(s)$ is real for $s \in i\mathbb{R}$. Then $\frac{3}{2\pi i}(\psi_R - \psi_P)(s)ds$ is a real measure on the imaginary axis. As $s \to \infty$, we have $\psi_R(s) - \psi_P(s) = 2 + \mathcal{O}(\frac{1}{s})$. Since $\psi_R - \psi_P$ never vanishes in $\overline{\mathbb{C}}$, this implies that $\frac{3}{2\pi i}(\psi_R - \psi_P)(s)ds$ is a positive measure on the imaginary axis, and so in particular on the interval $[-iy^*, iy^*]$. Hence all three parts of μ_Q in (2.14) are positive, so that μ_Q is a positive measure.

Now we want to calculate the total mass of μ_Q. If we separate in $\int d\mu_Q$ the contributions from ψ_P, ψ_Q, and ψ_R, we get

$$\int d\mu_Q = \frac{3}{2\pi i}\int_{z_1}^{z_2} \psi_P(s)ds + \frac{3}{2\pi i}\int_{z_2}^{z_3} \psi_Q(s)ds + \frac{3}{2\pi i}\int_{z_3}^{z_4} \psi_R(s)ds + \frac{3}{2\pi i}\int_{z_4}^{z_1} \psi_Q(s)ds.$$

Here $\int_{z_j}^{z_k}$ stands for integration along the path from z_j to z_k within Γ_Q. We deform the contour from z_1 to z_2 to the contour Γ_P. On Γ_P we have $(\psi_P)_+ = (\psi_Q)_-$. Thus

$$\frac{3}{2\pi i}\int_{z_1}^{z_2} \psi_P(s)ds = \frac{3}{2\pi i}\int_{\Gamma_P} (\psi_Q)_-(s)ds.$$

Similarly, we have

$$\frac{3}{2\pi i}\int_{z_3}^{z_4} \psi_R(s)ds = \frac{3}{2\pi i}\int_{\Gamma_R} (\psi_Q)_-(s)ds.$$

Therefore,

$$\int d\mu_Q = \frac{3}{2\pi i}\oint_{\gamma} \psi_Q(s)ds$$

where γ is the closed contour, consisting of the $-$sides of Γ_P and Γ_R and of the paths from z_2 to z_3 and from z_4 to z_1 within Γ_Q. Now we can use the residue theorem for the exterior domain of γ. The only contribution comes from infinity, and since $\psi_Q(s) = \frac{1}{3s} + \mathcal{O}(\frac{1}{s^2})$ as $s \to \infty$, see (3.3), we find $\int_{\Gamma_Q} d\mu_Q = 1$. Hence μ_Q is a probability measure on Γ_Q. □

4.7 The function g_Q

For g_Q we have properties that are analogous to the properties of g_P and g_R obtained in Lemmas 4.3 and 4.5.

Lemma 4.9. *For the derivative of* g_Q *we have*

$$g_Q'(z) = \begin{cases} 3\psi_Q(z), & z \in D_\infty, \\ 3\psi_P(z), & z \in D_P, \\ 3\psi_R(z), & z \in D_R. \end{cases} \tag{4.22}$$

Proof. We have

$$g_Q'(z) = \int \frac{1}{z-s} d\mu_Q(s).$$

As we did in the proof of Proposition 4.8 to evaluate $\int d\mu_Q(s)$, we can write the integral as an integral over the closed contour γ that consists of the $-$sides of Γ_P and Γ_R and of the paths from z_2 to z_3 and from z_4 to z_1 within Γ_Q. We get for $z \in D_\infty$,

$$g_Q'(z) = \frac{3}{2\pi i}\oint_{\gamma} \frac{\psi_Q(s)}{z-s} ds.$$

Since ψ_Q is analytic in D_∞ and vanishes at infinity, the only contribution comes from the pole at $s = z$ and the result is that $g_Q'(z) = 3\psi_Q(z)$ if $z \in D_\infty$. This gives (4.22) for $z \in D_\infty$. The expressions in the other two domains follow by analytic continuation. □

Lemma 4.10. g_Q *has the explicit representation*

$$g_Q(z) = 3zw - \log[w(w^2-1)] - 1 + \log\left(-\frac{1}{3}\right), \ \textit{with} \begin{cases} w = \psi_Q(z) & \textit{if } z \in D_\infty, \\ w = \psi_P(z) & \textit{if } z \in D_P, \\ w = \psi_R(z) & \textit{if } z \in D_R. \end{cases} \tag{4.23}$$

Proof. This is analogous to the proof of Lemma 4.5. □

We conclude this section with connections between g_P, g_Q and g_R that easily follow from the explicit expressions (4.9), (4.10), and (4.23).

Corollary 4.11. (a) *For $z \in D_\infty$, we have*

$$g_P(z) + g_Q(z) + g_R(z) = 3 \log z. \tag{4.24}$$

(b) *For $z \in D_P$, we have*

$$g_P(z) - g_Q(z) = 3z + \ell. \tag{4.25}$$

(c) *For $z \in D_R$, we have*

$$g_R(z) - g_Q(z) = -3z + \ell. \tag{4.26}$$

Proof. Adding the formulas (4.9), (4.10), and (4.23) with $z \in D_\infty$, we get

$$g_P(z) + g_Q(z) + g_R(z) = \\ 3z(w_1 + w_2 + w_3) - \log\left[w_1(w_1^2 - 1)w_2(w_2^2 - 1)w_3(w_3^2 - 1)\right] - 3 + \log\left(-\frac{4}{27}\right), \tag{4.27}$$

where $w_1 = \psi_P(z)$, $w_2 = \psi_Q(z)$, and $w_3 = \psi_R(z)$. Since w_1, w_2, and w_3 are the solutions of the cubic equation (2.4) we have

$$z(w - w_1)(w - w_2)(w - w_3) = zw^3 - w^2 - zw + \frac{1}{3}$$

for every $w \in \mathbb{C}$. Comparing coefficients of w^2, we get $z(w_1 + w_2 + w_3) = 1$, and taking $w = 0$, $w = 1$, and $w = -1$, we get $zw_1w_2w_3 = -\frac{1}{3}$, $z(w_1 - 1)(w_2 - 1)(w_3 - 1) = \frac{2}{3}$, and $z(w_1 + 1)(w_2 + 1)(w_3 + 1) = \frac{2}{3}$, respectively. Using this in (4.27), we arrive at (4.24).

For parts (b) and (c), we start with the equation (4.24) in D_∞. Making use of the jump relation (4.14) and (4.15) to go into D_P and D_R, we obtain (4.25) and (4.26). ☐

5 The Riemann–Hilbert problem and the first two transformations

Our asymptotic analysis is based on the Riemann–Hilbert problem for Y formulated in the introduction, see (1.8), (1.9). In this section we prove that the Riemann–Hilbert problem has a unique solution and that the solution is given in terms of the polynomials P_n, Q_n, and R_n, and the remainder E_n. We also do the first two transformation of the Riemann–Hilbert problem, which consist of a normalization of the problem at infinity, and a deformation of contours.

5.1 The Riemann–Hilbert problem

We show that the Riemann–Hilbert problem for Y has a solution in terms of the Hermite-Padé polynomials.

Theorem 5.1. *Let P_n, Q_n, R_n, and E_n be as above. Then the solution of the Riemann–Hilbert problem for Y (see the introduction) is unique and is given by*

$$Y(z) = \begin{pmatrix} p_{n+1,n-1,n}(3nz) & z^{-3n-2}q_{n+1,n-1,n}(3nz) & r_{n+1,n-1,n}(3nz) \\ P_n(z) & z^{-3n-2}Q_n(z) & R_n(z) \\ p_{n,n-1,n+1}(3nz) & z^{-3n-2}q_{n,n-1,n+1}(3nz) & r_{n,n-1,n+1}(3nz) \end{pmatrix}, \tag{5.1}$$

for z outside Γ, and

$$Y(z) = \begin{pmatrix} p_{n+1,n-1,n}(3nz) & z^{-3n-2}e_{n+1,n-1,n}(3nz) & r_{n+1,n-1,n}(3nz) \\ P_n(z) & z^{-3n-2}E_n(z) & R_n(z) \\ p_{n,n-1,n+1}(3nz) & z^{-3n-2}e_{n,n-1,n+1}(3nz) & r_{n,n-1,n+1}(3nz) \end{pmatrix}, \tag{5.2}$$

for z inside Γ. In the first rows of (5.1) and (5.2) we use the Hermite–Padé polynomials of indices $n+1, n-1, n$ normalized so that $p_{n+1,n-1,n}(3nz)$ is a monic polynomial, and in the third row we use the Hermite–Padé polynomials of indices $n, n-1, n+1$ normalized so that $r_{n,n-1,n+1}(3nz)$ is monic.

Proof. The given Y is analytic inside and outside the contour Γ. This is clear from (5.1) and (5.2), except perhaps for the second column of (5.2) which has a possible singularity at the origin. However, the singularity is removable since $e_{n_1,n_2,n_3}(z) = \mathcal{O}(z^{n_1+n_2+n_3+2})$ as $z \to 0$. The asymptotic condition (1.9) is satisfied because of the fact that $p_{n+1,n-1,n}(3nz)$, $Q_n(z)$, and $r_{n,n-1,n+1}(3nz)$ are monic polynomials.

The jump condition can easily be checked. For the first and third entries in the first row it reads

$$(Y_{11})_+ = (Y_{11})_-, \qquad (Y_{13})_+ = (Y_{13})_-,$$

which is indeed so since Y_{11} and Y_{13} are both polynomials. We also have

$$\begin{aligned} (Y_{12})_+(z) &= z^{-3n-2}e_{n+1,n-1,n}(3nz) \\ &= z^{-3n-2}\left(p_{n+1,n-1,n}(3nz)e^{-3nz} + q_{n+1,n-1,n}(3nz) + r_{n+1,n-1,n}(3nz)e^{3nz}\right) \\ &= z^{-3n-2}e^{-3nz}(Y_{11})_-(z) + (Y_{12})_-(z) + z^{-3n-2}e^{3nz}(Y_{13})_-(z), \end{aligned}$$

and this is the jump condition (1.8) for the second entry in the first row. The second and third rows are handled in the same way.

To prove uniqueness, we assume that $\widetilde{Y}$ is another solution of the Riemann–Hilbert problem. First observe that $\det Y$ is a scalar function which is analytic in $\mathbb{C} \setminus \Gamma$. Because of (1.8), we have that $(\det Y)_+(z) = (\det Y)_-(z)$ for $z \in \Gamma$, so that $\det Y$ has no jump, making $\det Y$ an entire function. For large z we have $\det Y(z) = 1 + \mathcal{O}(1/z)$ by (1.9), hence by Liouville's theorem $\det Y = 1$ everywhere. We can therefore consider $\widetilde{Y}Y^{-1}$, which is analytic in $\mathbb{C} \setminus \Gamma$. There is no jump on Γ since $(\widetilde{Y}Y^{-1})_+(z) = (\widetilde{Y}Y^{-1})_-(z)$ for every $z \in \Gamma$, hence $\widetilde{Y}Y^{-1}$ is entire (i.e., each entry is an entire function). For large z we have $\widetilde{Y}Y^{-1}(z) = I + \mathcal{O}(1/z)$, hence Liouville's theorem implies that $\widetilde{Y}Y^{-1}(z) = I$ for every z, and hence $\widetilde{Y}(z) = Y(z)$. ☐

Remark 5.2. A Riemann–Hilbert characterization for general Hermite–Padé polynomials was given in [39].

5.2 First transformation

We will use the functions g_P and g_R, and the constant $\ell = \log 2 - \pi i$ from Section 4.3 to transform the Riemann–Hilbert problem for Y to a Riemann–Hilbert problem for U, given by

$$U(z) = L^{-n-1} Y(z) \begin{pmatrix} e^{-(n+1)g_P(z)} & 0 & 0 \\ 0 & e^{(n+1)[g_P(z)+g_R(z)]} & 0 \\ 0 & 0 & e^{-(n+1)g_R(z)} \end{pmatrix} L^{n+1}, \tag{5.3}$$

where L is the constant diagonal matrix

$$L = \begin{pmatrix} e^{\ell/3} & 0 & 0 \\ 0 & e^{-2\ell/3} & 0 \\ 0 & 0 & e^{\ell/3} \end{pmatrix}. \tag{5.4}$$

For the contour Γ we take $\Gamma = \Gamma_P \cup \Gamma_R \cup \Gamma_U \cup \Gamma_L$, where Γ_U is a contour connecting z_4 to z_1 and lying in $D_{\infty,U}$, and Γ_L is a contour connecting z_2 to z_3 and lying in $D_{\infty,L}$. Then $\operatorname{Re}\varphi_P > 0$ and $\operatorname{Re}\varphi_R > 0$ on $\Gamma_U \cup \Gamma_L$ by Lemma 2.7.

We note that U is analytic on $\mathbb{C} \setminus \Gamma$, since $e^{g_P(z)}$ and $e^{g_R(z)}$ are analytic and single-valued on $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R)$ and $\Gamma_P \cup \Gamma_R \subset \Gamma$.

Since $g_P(z) = \log z + \mathcal{O}(1/z)$ as $z \to \infty$, we have $e^{(n+1)g_P(z)} = z^{n+1}[1+\mathcal{O}(1/z)]$ as $z \to \infty$. Similarly we also have $e^{(n+1)g_R(z)} = z^{n+1}[1+\mathcal{O}(1/z)]$ as $z \to \infty$. Hence

$$U(z) = I + \mathcal{O}\left(\frac{1}{z}\right), \qquad z \to \infty. \tag{5.5}$$

So U is normalized at infinity.

The jump relation for U needs to be worked out on the four pieces of the contour $\Gamma = \Gamma_P \cup \Gamma_R \cup \Gamma_U \cup \Gamma_L$.

For $z \in \Gamma_P$ we have

$$U_+(z) = U_-(z) \begin{pmatrix} e^{-(n+1)[g_{P+}(z)-g_{P-}(z)]} & z e^{3z} e^{(n+1)[-3\log z - 3z + g_{P+}(z) + g_{P-}(z) + g_R(z) - \ell]} & 0 \\ 0 & e^{(n+1)[g_{P+}(z)-g_{P-}(z)]} & 0 \\ 0 & z e^{-3z} e^{(n+1)[-3\log z + 3z + g_{P+}(z) + 2g_R(z) - \ell]} & 1 \end{pmatrix}. \tag{5.6}$$

Taking into account Lemma 4.6, we can simplify the jump (5.6) to

$$U_+(z) = U_-(z) \begin{pmatrix} e^{2(n+1)\varphi_{P+}(z)} & z e^{3z} & 0 \\ 0 & e^{2(n+1)\varphi_{P-}(z)} & 0 \\ 0 & z e^{-3z} e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}, \qquad z \in \Gamma_P.$$

Similarly, the jump on Γ_R is

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & z e^{3z} e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & e^{2(n+1)\varphi_{R-}(z)} & 0 \\ 0 & z e^{-3z} & e^{2(n+1)\varphi_{R+}(z)} \end{pmatrix}, \qquad z \in \Gamma_R.$$

On the parts Γ_U and Γ_L we have

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & z e^{3z} e^{(n+1)[-3\log z - 3z + 2g_P(z) + g_R(z) - \ell]} & 0 \\ 0 & 1 & 0 \\ 0 & z e^{-3z} e^{(n+1)[-3\log z + 3z + g_P(z) + 2g_R(z) - \ell]} & 1 \end{pmatrix}. \tag{5.7}$$

If we use Lemma 4.6 then (5.7) can be re-written as

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}, \qquad z \in \Gamma_U \cup \Gamma_L.$$

Summarizing, we have the following Riemann–Hilbert problem for U

1. U is analytic on $\mathbb{C} \setminus \Gamma$.
2. U satisfies the following jump relations

$$U_+(z) = U_-(z) \begin{pmatrix} e^{2(n+1)\varphi_{P+}(z)} & ze^{3z} & 0 \\ 0 & e^{2(n+1)\varphi_{P-}(z)} & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}, \qquad z \in \Gamma_P, \tag{5.8}$$

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & e^{2(n+1)\varphi_{R-}(z)} & 0 \\ 0 & ze^{-3z} & e^{2(n+1)\varphi_{R+}(z)} \end{pmatrix}, \qquad z \in \Gamma_R, \tag{5.9}$$

$$U_+(z) = U_-(z) \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}, \qquad z \in \Gamma_U \cup \Gamma_L. \tag{5.10}$$

3. $U(z) = I + \mathcal{O}\left(\frac{1}{z}\right)$ as $z \to \infty$.

In Figure 6 the shaded region is where $\operatorname{Re}\varphi_P$ is negative. For $\operatorname{Re}\varphi_R$ there is a similar picture, but reflected along the imaginary axis. The contours Γ_U and Γ_L are in the region where both $\operatorname{Re}\varphi_P$ and $\operatorname{Re}\varphi_R$ are positive. The jump matrix in (5.10) for U on the contours Γ_U and Γ_L is then the identity matrix I plus a matrix with entries that tend to zero exponentially fast as $n \to \infty$.

The factor $e^{-2(n+1)\varphi_R(z)}$ in the last row of the jump matrix in (5.8) tends exponentially fast to 0 because $\operatorname{Re}\varphi_R > 0$ on Γ_P, see Lemma 3.1. In a similar way the factor $e^{-2(n+1)\varphi_P(z)}$ in the first row of the jump matrix in (5.9) on Γ_R tends to 0 exponentially fast since $\operatorname{Re}\varphi_P > 0$ on Γ_R (see Lemma 3.1 and Figure 6). Furthermore $\varphi_{P+} = -\varphi_{P-}$ is purely imaginary on Γ_P because of (2.7) and $\varphi_{R+} = -\varphi_{R-}$ is purely imaginary on Γ_R because of (2.8), so that the diagonal elements of the jump matrices on Γ_P and Γ_R are oscillatory.

5.3 Deformation of contours

The jump matrix in (5.8) can be written as a product of four matrices

$$\begin{aligned} &\begin{pmatrix} e^{2(n+1)\varphi_{P+}(z)} & ze^{3z} & 0 \\ 0 & e^{2(n+1)\varphi_{P-}(z)} & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix} \\ &\qquad = \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_{P-}(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & ze^{3z} & 0 \\ -z^{-1}e^{-3z} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &\qquad\quad \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_{P+}(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}. \end{aligned} \tag{5.11}$$

Instead of jumping over Γ_P in one jump, we will make four smaller jumps, and rather than jumping over one contour, we jump over four contours, and each contour deals with one of the matrices in the product (5.11). We will open up a lens around Γ_P and introduce an extra contour for the last factor of the matrix. The lens consists of two contours $\Gamma_{P-} \cup \Gamma_{P+}$ connecting z_1 and z_2, such that Γ_{P-} is on the minus side of Γ_P and Γ_{P+} is on the plus side of Γ_P, but still inside the region where $\operatorname{Re}\varphi_R > 0$ and $\operatorname{Re}\varphi_P < 0$. The contour Γ_{P++} stays away from z_1 and z_2. It starts at a point on the upper contour Γ_U and connects with a point on the lower contour Γ_L, so that it lies inside the region where $\operatorname{Re}\varphi_R > 0$, (so it stays to the left of Γ_R^*), and does not intersect Γ_{P+}. The subarc of Γ_U between z_1 and the starting point of Γ_{P++} we call Γ_{U^p}, and the subarc of Γ_L from z_2 to the endpoint of Γ_{P++} is called Γ_{L^p}.

These contours are drawn in Figure 7.



Figure 7: Deformation of contours around Γ_P. The dotted line is Γ_R^*

In a similar way the jump (5.9) over Γ_R has a factorization

$$\begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & e^{2(n+1)\varphi_{R-}(z)} & 0 \\ 0 & ze^{3z} & e^{2(n+1)\varphi_{R+}(z)} \end{pmatrix}$$
$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z^{-1}e^{3z}e^{2(n+1)\varphi_{R-}(z)} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -z^{-1}e^{3z} \\ 0 & ze^{-3z} & 0 \end{pmatrix}$$
$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z^{-1}e^{3z}e^{2(n+1)\varphi_{R+}(z)} \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (5.12)$$

We now open up contours Γ_{R-}, Γ_{R^+}, and $\Gamma_{R^{++}}$ around Γ_R, which are mirror images of the contours around Γ_P. The mirror images of Γ_{U^p} and Γ_{L^p} are denoted by Γ_{U^r} and Γ_{L^r}, respectively. The remaining middle parts of Γ_U and Γ_L are denoted by Γ_{U^m} and Γ_{L^m}, respectively.



Figure 8: Contours of the RHP for T

All these contours are shown in Figure 8. All together there are 14 contours, and they determine 8 regions in the plane. The second transformation $U \mapsto T$ will be defined in each of these regions separately. We define T as follows.

We take

$$T(z) = U(z), \tag{5.13}$$

for z in the unbounded region, and in the middle region bounded by $\Gamma_{P^{++}}$, $\Gamma_{R^{++}}$, Γ_{U^m}, and Γ_{L^m}. In the three regions near Γ_P we put

$$T(z) = \begin{cases} U(z)V_{P^-}(z), & \text{for } z \text{ in the region bounded by } \Gamma_{P^-} \text{ and } \Gamma_P, \\ U(z)V_{P^{++}}^{-1}(z)V_{P^+}^{-1}(z), & \text{for } z \text{ in the region bounded by } \Gamma_P \text{ and } \Gamma_{P^+}, \\ U(z)V_{P^{++}}^{-1}(z), & \text{for } z \text{ in the region bounded by } \Gamma_{P^+}, \Gamma_{P^{++}}, \Gamma_{U^p}, \text{ and } \Gamma_{L^p}, \end{cases} \tag{5.14}$$

where

$$V_{P^-}(z) = \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.15}$$

$$V_{P^+}(z) = \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.16}$$

$$V_{P^{++}}(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}. \tag{5.17}$$

In the regions near Γ_R we put similarly

$$T(z) = \begin{cases} U(z)V_{R^-}(z), & \text{for } z \text{ in the region bounded by } \Gamma_{R^-} \text{ and } \Gamma_R, \\ U(z)V_{R^{++}}^{-1}(z)V_{R^+}^{-1}(z), & \text{for } z \text{ in the region bounded by } \Gamma_R \text{ and } \Gamma_{R^+}, \\ U(z)V_{R^{++}}^{-1}(z), & \text{for } z \text{ in the region bounded by } \Gamma_{R^+}, \Gamma_{R^{++}}, \Gamma_{U^r}, \text{ and } \Gamma_{L^r}, \end{cases} \tag{5.18}$$

where

$$V_{R^-}(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z^{-1}e^{3z}e^{2(n+1)\varphi_R(z)} \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.19}$$

$$V_{R^+}(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & z^{-1}e^{3z}e^{2(n+1)\varphi_R(z)} \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.20}$$

$$V_{R^{++}}(z) = \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{5.21}$$

Then we have the following Riemann–Hilbert problem for T.

1. T is analytic in each of the 8 regions,
2. T has a jump on each of the 14 contours

   $$T_+(z) = T_-(z)V_s(z), \qquad z \in \Gamma_s,$$

   where s stand for any of the symbols P, P^-, P^+, P^{++}, R, R^-, R^+, R^{++}, U^p, U^m, U^r, L^p, L^m, or L^r. The matrices V_{P^-}, V_{P^+}, $V_{P^{++}}$, V_{R^-}, V_{R^+}, and $V_{R^{++}}$ have already been defined above. The other jump matrices are

   $$V_P(z) = \begin{pmatrix} 0 & ze^{3z} & 0 \\ -z^{-1}e^{-3z} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.22}$$

   $$V_R(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -z^{-1}e^{3z} \\ 0 & ze^{-3z} & 0 \end{pmatrix}, \tag{5.23}$$

   $$V_{U^p}(z) = V_{L^p}(z) = \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{5.24}$$

   $$V_{U^r}(z) = V_{L^r}(z) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}, \tag{5.25}$$

   $$V_{U^m}(z) = V_{L^m}(z) = \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}. \tag{5.26}$$

3. $T(z) = I + \mathcal{O}\left(\frac{1}{z}\right)$ as $z \to \infty$.

Observe that all jumps, except for the jumps on Γ_P and Γ_R, tend to the identity matrix exponentially fast as $n \to \infty$. Hence we expect that the dominating contributions are the jumps V_P on Γ_P and V_R on Γ_R.

6 Construction of parametrices and final transformation

6.1 Parametrix for the exterior region

We will now solve a Riemann–Hilbert problem for a matrix valued function N on the contours $\Gamma_P \cup \Gamma_R$ (see Figure 9) which, in view of what was said at the end of the previous section, is expected to describe the main contribution of the Riemann–Hilbert problem of T.



Figure 9: Contours of the RHP for N

We look for $N : \mathbb{C} \setminus (\Gamma_P \cup \Gamma_R) \to \mathbb{C}^{3\times 3}$ satisfying

1. N is analytic in $\mathbb{C} \setminus (\Gamma_P \cup \Gamma_R)$.
2. N has jumps on Γ_P and Γ_R given by
$$N_+(z) = N_-(z) \begin{pmatrix} 0 & ze^{3z} & 0 \\ -z^{-1}e^{-3z} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad z \in \Gamma_P, \tag{6.1}$$
and
$$N_+(z) = N_-(z) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -z^{-1}e^{3z} \\ 0 & ze^{-3z} & 0 \end{pmatrix}, \qquad z \in \Gamma_R. \tag{6.2}$$
3. $N(z) = I + \mathcal{O}\left(\frac{1}{z}\right)$ as $z \to \infty$.

Proposition 6.1. *A solution of the Riemann–Hilbert problem for N is given by*

$$N(z) = \begin{pmatrix} F_1(\psi_P(z)) & F_1(\psi_Q(z)) & F_1(\psi_R(z)) \\ F_2(\psi_P(z)) & F_2(\psi_Q(z)) & F_2(\psi_R(z)) \\ F_3(\psi_P(z)) & F_3(\psi_Q(z)) & F_3(\psi_R(z)) \end{pmatrix}, \tag{6.3}$$

where

$$F_1(w) = \frac{-w(w-1)G(w)}{\sqrt{3w^4+1}}, \tag{6.4}$$

$$F_2(w) = \frac{3(w^2-1)G(w)}{\sqrt{3w^4+1}}, \tag{6.5}$$

$$F_3(w) = \frac{w(w+1)G(w)}{\sqrt{3w^4+1}}, \tag{6.6}$$

with $\sqrt{3w^4+1}$ defined and analytic in $\mathbb{C} \setminus (\psi_{P+}(\Gamma_P) \cup \psi_{R+}(\Gamma_R))$, and such that it is positive for large positive w. The function G is defined by

$$G(w) = \begin{cases} w e^{-\frac{(w+1)(2w-1)}{w(w-1)}} & \text{for } w \in \psi(\mathcal{R}_P), \\ \left(\frac{w^2-1/3}{w^2-1}\right) e^{-\frac{2w^2}{w^2-1}} & \text{for } w \in \psi(\mathcal{R}_Q), \\ w e^{-\frac{(w-1)(2w+1)}{w(w+1)}} & \text{for } w \in \psi(\mathcal{R}_R). \end{cases} \tag{6.7}$$

Proof. Let us consider the first row (N_{11}, N_{12}, N_{13}) of N. From (6.1) we get the following jumps on Γ_P

$$\begin{cases} (N_{11})_+(z) = -z^{-1}e^{-3z}(N_{12})_-(z), \\ (N_{12})_+(z) = ze^{3z}(N_{11})_-(z), \\ (N_{13})_+(z) = (N_{13})_-(z), \end{cases} \qquad z \in \Gamma_P, \tag{6.8}$$

and from (6.2) the following jumps on Γ_R

$$\begin{cases} (N_{11})_+(z) = (N_{11})_-(z), \\ (N_{12})_+(z) = ze^{-3z}(N_{13})_-(z), \\ (N_{13})_+(z) = -z^{-1}e^{3z}(N_{12})_-(z), \end{cases} \qquad z \in \Gamma_R. \tag{6.9}$$

Clearly N_{11} is analytic on Γ_R and N_{13} is analytic on Γ_P. So we can (and do) see N_{11} as a function on the sheet $\mathcal{R}_P$ of the Riemann surface $\mathcal{R}$, N_{12} as a function on $\mathcal{R}_Q$ and N_{13} as a function on $\mathcal{R}_R$. Then we transform the problem from $\mathcal{R}$ with the variable z, to the complex w-plane, via the mapping $\psi : \mathcal{R} \to \mathbb{C}$. The variables z and w are connected by (2.3). The images of the three sheets, the images of the branch points z_1, z_2, z_3, z_4, and the images of the cuts Γ_P and Γ_R are shown in Figure 3.

Note that the images of Γ_P under the mappings ψ_{P+} and ψ_{P-} (positive and negative boundary values of ψ_P on Γ_P) give two arcs from w_1 to w_2. They are oriented as shown in Figure 3. The orientation corresponds to the orientation of Γ_P. Together the arcs make up a simple closed loop around -1.

Similar remarks hold for the two images $\psi_{R+}(\Gamma_R)$ and $\psi_{R-}(\Gamma_R)$ of Γ_R. They make up a simple closed loop around 1.

Now we transplant the (as yet unknown) functions N_{11}, N_{12}, and N_{13} from the Riemann surface to the w-plane, by defining F_1 as follows.

$$F_1(w) = \begin{cases} N_{11}\left(\frac{w^2-\frac{1}{3}}{w(w^2-1)}\right), & w \in \psi(\mathcal{R}_P), \\ N_{12}\left(\frac{w^2-\frac{1}{3}}{w(w^2-1)}\right), & w \in \psi(\mathcal{R}_Q), \\ N_{13}\left(\frac{w^2-\frac{1}{3}}{w(w^2-1)}\right), & w \in \psi(\mathcal{R}_R). \end{cases} \tag{6.10}$$

Then F_1 is analytic in $\mathbb{C} \setminus (\psi_{P\pm}(\Gamma_P) \cup \psi_{R\pm}(\Gamma_R))$. The jumps that F_1 should satisfy can be determined from (6.8)–(6.9) and are given by

$$\begin{cases} F_{1+}(w) = ze^{3z}F_{1-}(w), & w \in \psi_{P-}(\Gamma_P), \\ F_{1+}(w) = -z^{-1}e^{-3z}F_{1-}(w), & w \in \psi_{P+}(\Gamma_P), \\ F_{1+}(w) = -z^{-1}e^{3z}F_{1-}(w), & w \in \psi_{R+}(\Gamma_R), \\ F_{1+}(w) = ze^{-3z}F_{1-}(w), & w \in \psi_{R-}(\Gamma_R), \end{cases} \tag{6.11}$$

where $z = z(w) = \frac{w^2-\frac{1}{3}}{w(w^2-1)}$.

The asymptotic condition on N implies that $N_{11}(z) \to 1$, $N_{12}(z) \to 0$, $N_{13}(z) \to 0$ as $z \to \infty$. For F_1, this means that

$$F_1(-1) = 1, \qquad F_1(0) = 0, \qquad F_1(1) = 0. \tag{6.12}$$

We also want $F_1(w)$ to have a finite limit as $w \to \infty$, since $w = \infty$ corresponds to $z = 0$ on the Q-sheet.

We now seek F_1 in the form

$$F_1(w) = \frac{-w(w-1)G(w)}{\sqrt{3w^4+1}}. \tag{6.13}$$

Then G should be analytic in $\mathbb{C} \setminus (\psi_{P\pm}(\Gamma_P) \cup \psi_{R\pm}(\Gamma_R))$ with jumps

$$\begin{cases} G_+(w) = ze^{3z}G_-(w), & w \in \psi_{P-}(\Gamma_P), \\ G_+(w) = z^{-1}e^{-3z}G_-(w), & w \in \psi_{P+}(\Gamma_P), \\ G_+(w) = z^{-1}e^{3z}G_-(w), & w \in \psi_{R+}(\Gamma_R), \\ G_+(w) = ze^{-3z}G_-(w), & w \in \psi_{R-}(\Gamma_R), \end{cases} \tag{6.14}$$

with $z = z(w)$. The normalization for G is

$$G(-1) = -1. \tag{6.15}$$

It is straightforward to check that G given by (6.7) indeed satisfies (6.14) and (6.15). Then by (6.13) it follows that F_1 has the correct jumps (6.11) and normalization (6.12). Then from (6.10) we recover N_{11}, N_{12}, and N_{13} in terms of F_1 by

$$N_{11}(z) = F_1(\psi_P(z)), \quad N_{12}(z) = F_1(\psi_Q(z)), \quad N_{13}(z) = F_1(\psi_R(z)).$$

Then the jumps (6.8) and (6.9) are satisfied, and in addition the normalization at infinity is correct. So we have found the first row of N.

The proof for the second and third rows is similar. The only difference is that we have a different normalization at infinity, which leads to the construction of functions F_2 and F_3 that satisfy the same jumps (6.11) as F_1, but are normalized by

$$F_2(-1) = 0, \qquad F_2(0) = 1, \qquad F_2(1) = 0,$$

and

$$F_3(-1) = 0, \qquad F_3(0) = 0, \qquad F_3(1) = 1.$$

Similar calculations then lead to the formulas (6.5) and (6.6) with the same function G. □

We remark that the entries of N have fourth root singularities at the branch points z_k ($k = 1, \ldots, 4$). More precisely,

Lemma 6.2. *The entries of N behave as follows near the branch points. As $z \to z_j$ with $j = 1, 2$, we have*

$$\begin{cases} N_{k1}(z) & = \mathcal{O}\left(|z-z_j|^{-1/4}\right), \\ N_{k2}(z) & = \mathcal{O}\left(|z-z_j|^{-1/4}\right), \\ N_{k3}(z) & = \mathcal{O}\left(1\right), \end{cases} \qquad k = 1, 2, 3, \tag{6.16}$$

and as $z \to z_j$ with $j = 3, 4$, we have

$$\begin{cases} N_{k1}(z) & = \mathcal{O}\left(1\right), \\ N_{k2}(z) & = \mathcal{O}\left(|z-z_j|^{-1/4}\right), \\ N_{k3}(z) & = \mathcal{O}\left(|z-z_j|^{-1/4}\right), \end{cases} \qquad k = 1, 2, 3. \tag{6.17}$$

Proof. Since w_1 is a non-degenerate critical point of the mapping $z = z(w)$, we have for the inverse $w = \psi_P(z)$ as $z \to z_1 = z(w_1)$,

$$\psi_P(z) = w_1 + c(z-z_1)^{1/2} + \mathcal{O}\left(z - z_1\right) \tag{6.18}$$

where c is a non-zero constant. Since w_1 is a simple root of $3w^4 + 1$, it then follows that

$$\sqrt{3\psi_P^4(z) + 1} = c_2(z-z_1)^{1/4} + \mathcal{O}\left(|z-z_1|^{3/4}\right), \qquad z \to z_1, \tag{6.19}$$

with $c_2 \neq 0$. Since the numerators of F_1, F_2, F_3 as given by (6.4)–(6.6), do not vanish for $w = w_1$, we find

$$N_{k1}(z) = F_k(\psi_P(z)) = \mathcal{O}\left(|z-z_1|^{-1/4}\right), \qquad k = 1, 2, 3,$$

as $z \to z_1$. In a similar way we find that $N_{k2}(z) = \mathcal{O}\left(|z-z_1|^{-1/4}\right)$ as $z \to z_1$. The entries in the third column of N are analytic at z_1, since ψ_R is analytic at z_1 and the functions F_1, F_2, F_3 are analytic at $\psi_R(z_1)$. This proves (6.16) for $j = 1$.

The behavior near the other branch points follows in a similar way. □

Remark 6.3. It will be useful to have another representation for the entries in the second row of N. They are

$$N_{21}(z) = \frac{2e^{-g_P(z)}}{\sqrt{3\psi_P^4(z)+1}}, \quad N_{22}(z) = -\frac{e^{g_P(z)+g_R(z)}}{z^2\sqrt{3\psi_Q^4(z)+1}}, \quad N_{23}(z) = \frac{2e^{-g_R(z)}}{\sqrt{3\psi_R^4(z)+1}}. \tag{6.20}$$

It may be checked directly that these functions have the right asymptotics as $z \to \infty$, and satisfy the correct jump relations on Γ_P and Γ_R. They also satisfy the $\mathcal{O}$-conditions of Lemma 6.2.

The Riemann–Hilbert problem for T is now very close to the Riemann–Hilbert problem for N because the jumps for T and N on the contours Γ_P and Γ_R are the same and the jumps for T on the other contours tend to the identity matrix as $n \to \infty$, uniformly away from the branch points. So we expect that T behaves like N as $n \to \infty$ away from the branch points. However, in order to justify this we need to analyze the Riemann–Hilbert problem in more detail near the branch points.

6.2 Parametrices near the branch points

We introduce four new contours Γ_1, Γ_2, Γ_3, Γ_4 which are small circles of radius δ centered at the branch points, as indicated in Figure 10. We choose δ small enough so that the circles do not intersect the curves Γ_{P++} and Γ_{R++}. The circles are oriented clockwise. Inside each of these contours we will solve the Riemann–Hilbert problem for T exactly. The analysis is similar for the four branch points, and we will only work out the analysis near z_1 in detail.



Figure 10: Small circles Γ_1, Γ_2, Γ_3, Γ_4 around the branch points

Zooming in near the branch point z_1 gives a Riemann–Hilbert problem with five contours $\Gamma_{P_1}, \Gamma_P, \Gamma_{P+}, \Gamma_U, \Gamma_1$, as shown in Figure 11.



Figure 11: The contours near the branch point z_1

The jumps on these contours are

$$\begin{aligned} V_{P^-}(z) = V_{P^+}(z) &= \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ V_P &= \begin{pmatrix} 0 & ze^{3z} & 0 \\ -z^{-1}e^{-3z} & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ V_{U^p}(z) &= \begin{pmatrix} 1 & ze^{3z}e^{-2(n+1)\varphi_P(z)} & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

We look for a 3×3 matrix valued function $M^{(1)}$ defined within the disk Δ_1 surrounded by Γ_1, such that

1. $M^{(1)}$ is analytic in $\Delta_1 \setminus (\Gamma_P \cup \Gamma_{P^-} \cup \Gamma_{P^+} \cup \Gamma_{U^p})$,
2. $M^{(1)}$ has the jumps

   $$M_+^{(1)}(z) = M_-^{(1)}(z)V_s(z), \qquad z \in \Gamma_s,$$

   where s stands for any of the symbols P, P^-, P^+, and U^p.
3. On Γ_1 we have that $M^{(1)}$ matches N in the sense that

   $$M^{(1)}(z) = \left(I + \mathcal{O}\left(\frac{1}{n}\right)\right) N(z)$$

   uniformly for $z \in \Gamma_1$.

The construction of $M^{(1)}$ is basically a 2×2 problem, since the jump matrices are non-trivial only in the 2×2 left upper block. The jumps have a standard form and a local parametrix can be built out of Airy functions. To be more precise, we will find $M^{(1)}$ in the form

$$M^{(1)}(z) = E^{(1)}(z)\Psi^{(1)}((n+1)^{2/3}f_1(z)) \begin{pmatrix} z^{-1/2}e^{-3z/2}e^{(n+1)\varphi_P(z)} & 0 & 0 \\ 0 & z^{1/2}e^{3z/2}e^{-(n+1)\varphi_P(z)} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \tag{6.21}$$

Here we take the branch of φ_P which is zero at z_1. So for $z \in \Delta_1$, we take

$$\varphi_P(z) = \frac{3}{2}\int_{z_1}^{z} (\psi_Q - \psi_P)(s)ds$$

with integration from z_1 to z in $\Delta_1 \setminus \Gamma_P$. Recall that φ_P is defined modulo πi, and that so far the precise branch did not matter, since we only had expressions like $e^{2(n+1)\varphi_P(z)}$. Since

$$\psi_P(s) = w_1 + c(s-z_1)^{1/2} + \mathcal{O}(s-z_1), \quad \psi_Q(s) = w_1 - c(s-z_1)^{1/2} + \mathcal{O}(s-z_1),$$

as $s \to z_1$, with the same non-zero constant c_1, see also (6.18), we then have from this definition of φ_P that

$$\varphi_P(z) = (z-z_1)^{3/2}h(z), \qquad z \in \Delta_1 \setminus \Gamma_P,$$

with h analytic and without zeros in Δ_1. The function $(z-z_1)^{3/2}$ is defined with a branch cut along Γ_P.

The function $f_1(z)$ is defined such that

$$\varphi_P(z) = \frac{2}{3}\left[f_1(z)\right]^{3/2}.$$

It is a conformal map from Δ_1 onto a convex neighborhood of 0. (We may have to shrink Δ_1, if necessary.) It maps Γ_P onto a part of the negative real axis. We still have some freedom in the choice of Γ_{P-}, Γ_{P+} and Γ_{U^p}. We take Γ_{U^p} so that f_1 maps Γ_{U^p} to a part of the positive real line. This means that Γ_{U^p} is an analytic continuation of Γ_P. Γ_{P-} and Γ_{P+} are chosen so that they are mapped by f_1 onto rays in the complex s-plane. We denote the images of Γ_P, Γ_{P-}, Γ_{P+}, and Γ_{U^p}, by Σ_P, Σ_{P-}, Σ_{P+} and Σ_{U^p}. These contours are shown in Figure 12. On these contours we use the constant jump matrices

$$\widehat{V}_{P-} = \widehat{V}_{P+} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

on Σ_{P-} and Σ_{P+},

$$\widehat{V}_P = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

on Σ_P, and

$$\widehat{V}_{U^p} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

on Σ_{U^p}. This Riemann–Hilbert problem is well-known when we deal with 2×2 matrices (see, e.g., [12, §7.6], [13], [14, p. 1522]). Note however that the arrows are pointing in the opposite direction compared to Figures 7.18–7.20 in [12], so that we need to modify the solution a little. The solution $\Psi^{(1)}$ uses the Airy function $\operatorname{Ai}(z)$, which is the unique solution of $y''(z) = zy(z)$



Figure 12: The contours for $\Psi^{(1)}$

with asymptotics as $z \to \infty$ given by

$$\operatorname{Ai}(z) = \frac{1}{2\sqrt{\pi}} z^{-1/4} e^{-\frac{2}{3}z^{3/2}} \left(1 + \mathcal{O}\left(\frac{1}{z^{3/2}}\right)\right), \tag{6.22}$$

$$\operatorname{Ai}'(z) = \frac{-1}{2\sqrt{\pi}} z^{1/4} e^{-\frac{2}{3}z^{3/2}} \left(1 + \mathcal{O}\left(\frac{1}{z^{3/2}}\right)\right), \tag{6.23}$$

which holds for $|\arg z| < \pi$, where $z^{1/4}$ and $z^{3/2}$ are defined with principal branch (i.e., with a cut along the negative real axis). We will define $\Psi^{(1)}$ by

$$\Psi^{(1)}(s) = \begin{pmatrix} \operatorname{Ai}(s) & -\operatorname{Ai}(\omega_3^2 s) & 0 \\ \operatorname{Ai}'(s) & -\omega_3^2 \operatorname{Ai}'(\omega_3^2 s) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\pi/6} & 0 & 0 \\ 0 & e^{i\pi/6} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad s \in I,$$

$$\Psi^{(1)}(s) = \begin{pmatrix} \operatorname{Ai}(s) & -\operatorname{Ai}(\omega_3^2 s) & 0 \\ \operatorname{Ai}'(s) & -\omega_3^2 \operatorname{Ai}'(\omega_3^2 s) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\pi/6} & 0 & 0 \\ 0 & e^{i\pi/6} & 0 \\ 0 & 0 & 1 \end{pmatrix} \widehat{V}_{P-}, \qquad s \in II,$$

$$\Psi^{(1)}(s) = \begin{pmatrix} \operatorname{Ai}(s) & \omega_3^2 \operatorname{Ai}(\omega_3 s) & 0 \\ \operatorname{Ai}'(s) & \operatorname{Ai}'(\omega_3 s) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\pi/6} & 0 & 0 \\ 0 & e^{i\pi/6} & 0 \\ 0 & 0 & 1 \end{pmatrix} \widehat{V}_{P+}^{-1}, \qquad s \in III,$$

$$\Psi^{(1)}(s) = \begin{pmatrix} \operatorname{Ai}(s) & \omega_3^2 \operatorname{Ai}(\omega_3 s) & 0 \\ \operatorname{Ai}'(s) & \operatorname{Ai}'(\omega_3 s) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{-i\pi/6} & 0 & 0 \\ 0 & e^{i\pi/6} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad s \in IV,$$

where $\omega_3 = e^{2\pi i/3}$ is a primitive third root of unity.

With the above definitions of $\Psi^{(1)}$ and f_1 it may then be shown that for any analytic prefactor $E^{(1)}$ the matrix $M^{(1)}$ defined by (6.21) satisfies the jump conditions on Γ_s, where s is any of the symbols P, P^-, P^+, and U^p. The extra factor $E^{(1)}$ has to be chosen in such a way that $M^{(1)}$ satisfies the matching condition on Γ_1 as well.

On the part of Γ_1 that lies in $(f_1)^{-1}(I)$ (the arc between Γ_{P-} and Γ_{U^p}) the asymptotic expansions for the Airy functions (6.22)–(6.23) give

$$\begin{aligned}
\mathrm{Ai}((n+1)^{2/3} f_1(z)) &= \frac{1}{2\sqrt{\pi}}(n+1)^{-1/6}\,(f_1(z))^{-\frac{1}{4}}\,e^{-(n+1)\varphi_P(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right),\\
\mathrm{Ai}((n+1)^{2/3}\omega_3^2 f_1(z)) &= \frac{1}{2\sqrt{\pi}}(n+1)^{-1/6}\,(f_1(z))^{-\frac{1}{4}}\,e^{i\pi/6}e^{(n+1)\varphi_P(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right),\\
\mathrm{Ai}'((n+1)^{2/3} f_1(z)) &= \frac{-1}{2\sqrt{\pi}}(n+1)^{1/6}\,(f_1(z))^{\frac{1}{4}}\,e^{-(n+1)\varphi_P(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right),\\
\mathrm{Ai}'((n+1)^{2/3}\omega_3^2 f_1(z)) &= \frac{-1}{2\sqrt{\pi}}(n+1)^{1/6}\,(f_1(z))^{\frac{1}{4}}\,e^{-i\pi/6}e^{(n+1)\varphi_P(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right).
\end{aligned}$$

Here the fourth root in $(f_1(z))^{\frac{1}{4}}$ is defined with a cut along Γ_P. This means that on this part of Γ_1 we have

$$\begin{aligned}
M^{(1)}(z) = \frac{1}{2\sqrt{\pi}}E^{(1)}(z)&\begin{pmatrix}((n+1)^{2/3}f_1(z))^{-\frac{1}{4}} & 0 & 0\\ 0 & ((n+1)^{2/3}f_1(z))^{\frac{1}{4}} & 0\\ 0 & 0 & 1\end{pmatrix}\\
&\begin{pmatrix}e^{-i\pi/6} & -e^{i\pi/3} & 0\\ -e^{-i\pi/6} & -e^{i\pi/3} & 0\\ 0 & 0 & 1\end{pmatrix}\begin{pmatrix}z^{-1/2}e^{-3z/2} & 0 & 0\\ 0 & z^{1/2}e^{3z/2} & 0\\ 0 & 0 & 1\end{pmatrix}\left(I+\mathcal{O}\left(\frac{1}{n}\right)\right). \quad (6.24)
\end{aligned}$$

In order to have the matching condition we therefore choose

$$\begin{aligned}
E^{(1)}(z) = 2\sqrt{\pi}N(z)&\begin{pmatrix}z^{1/2}e^{3z/2} & 0 & 0\\ 0 & z^{-1/2}e^{-3z/2} & 0\\ 0 & 0 & 1\end{pmatrix}\\
&\begin{pmatrix}e^{-i\pi/6} & -e^{i\pi/3} & 0\\ -e^{-i\pi/6} & -e^{i\pi/3} & 0\\ 0 & 0 & 1\end{pmatrix}^{-1}\begin{pmatrix}((n+1)^{2/3}f_1(z))^{\frac{1}{4}} & 0 & 0\\ 0 & ((n+1)^{2/3}f_1(z))^{-\frac{1}{4}} & 0\\ 0 & 0 & 1\end{pmatrix}\\
&= \sqrt{\pi}e^{i\pi/6}N(z)\begin{pmatrix}z^{1/2}e^{3z/2} & 0 & 0\\ 0 & z^{-1/2}e^{-3z/2} & 0\\ 0 & 0 & 1\end{pmatrix}\\
&\begin{pmatrix}1 & -1 & 0\\ i & i & 0\\ 0 & 0 & 2e^{-i\pi/6}\end{pmatrix}\begin{pmatrix}((n+1)^{2/3}f_1(z))^{\frac{1}{4}} & 0 & 0\\ 0 & ((n+1)^{2/3}f_1(z))^{-\frac{1}{4}} & 0\\ 0 & 0 & 1\end{pmatrix}. \quad (6.25)
\end{aligned}$$

With this choice of $E^{(1)}$ it is then easy to see that the matching condition holds on the part of Γ_1 in the region bounded by Γ_{P^-} and Γ_{U^p}. A similar analysis for the other parts of Γ_1 shows that the same $E^{(1)}$ also works in the other regions.

From (6.25) it is easy to see that $E^{(1)}$ is analytic in $\Delta_1 \setminus \Gamma_P$. On Γ_P, both N and $(f_1)^{\frac{1}{4}}$ have a jump. N has the jump (6.1) and $(f_1)^{\frac{1}{4}}$ satisfies $(f_1)_+^{\frac{1}{4}} = -i\,(f_1)_-^{\frac{1}{4}}$. Straightforward calculations then show that $E_+^{(1)} = E_-^{(1)}$ on Γ_P, so that $E^{(1)}$ is analytic across Γ_P. From (6.25) and the fact that the entries of N have at most fourth root singularities at z_1, see (6.16), we see that the entries of $E^{(1)}$ have at most a square root singularity at z_1. Since $E^{(1)}$ is analytic in $\Delta_1 \setminus \{z_1\}$, the singularity at z_1 is removable, and this proves that $E^{(1)}$ is analytic in the full Δ_1. This completes the construction of the parametrix $M^{(1)}$ in the neighborhood Δ_1 of z_1.

From the definitions (6.21) and (6.25) it is clear that

$$M^{(1)} = N \begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

where $*$ denotes an unspecified entry. This means that the third column of $M^{(1)}$ agrees with the third column of N. We will use this in what follows.

In a similar way, we can construct parametrices $M^{(2)}$, $M^{(3)}$, and $M^{(4)}$ near the other branch points z_2, z_3, z_4. The third column of $M^{(2)}$ agrees with the third column of N, and the first columns of $M^{(3)}$ and $M^{(4)}$ agree with the first column of N.

6.3 Third transformation

We now introduce the final matrix

$$S(z) = \begin{cases} T(z)\left(N(z)\right)^{-1}, & z \text{ outside } \Gamma_1, \Gamma_2, \Gamma_3, \Gamma_4, \\ T(z)\left(M^{(j)}(z)\right)^{-1}, & z \text{ inside } \Gamma_j,\ j = 1, 2, 3, 4. \end{cases} \tag{6.26}$$

Inside each Γ_j the matrices T and $M^{(j)}$ have the same jumps, hence S has no jumps inside Γ_j. Outside the Γ_j the matrices T and N have the same jump matrices on Γ_P and Γ_R. Hence S has no jump on Γ_P and Γ_R. This means that S solves a Riemann–Hilbert problem on the system of curves shown in Figure 13.

S is analytic outside the above system of contours and it is normalized at infinity

$$S(z) = I + \mathcal{O}\left(\frac{1}{z}\right), \qquad z \to \infty. \tag{6.27}$$

Theorem 6.4. *The matrix $S(z)$ has the behavior*

$$S(z) = I + \mathcal{O}\left(\frac{1}{n}\right), \tag{6.28}$$

uniformly on $\mathbb{C} \setminus \Sigma_S$, where Σ_S are the contours in Figure 13.

Proof. The jumps on all of the contours are uniformly of the form $I + \mathcal{O}(e^{-cn})$ with some fixed $c > 0$, except for the jumps on the circles Γ_j where we have

$$S_+(z) = S_-(z) M^{(j)}(z) N^{-1}(z), \qquad z \in \Gamma_j.$$

Because of the matching condition we have

$$M^{(j)}(z) N^{-1}(z) = I + \mathcal{O}\left(\frac{1}{n}\right)$$

uniformly for $z \in \Gamma_j$. Hence $S(z)$ solves a Riemann–Hilbert problem, normalized at ∞ with jumps close to the identity matrix up to $\mathcal{O}(1/n)$, uniformly on the contours Σ_S. We can then use arguments as those leading to Theorem 7.171 in [12] to obtain (6.28), see also [13, 14, 21]. □



Figure 13: Contours of the RHP for S

Using the asymptotic expansion for the Airy function and its derivative, it is possible to obtain a full asymptotic expansion for the jump matrix on the contours Γ_j in powers of $1/n$. This in turn yields a full asymptotic expansion for S

$$S(z) \sim I + \sum_{k=1}^{\infty} \frac{S_k(z)}{n^k} \tag{6.29}$$

as in Theorem 7.10 of [14]. We will not use the full expansion, since (6.28) is enough for the asymptotic results as stated in Section 2. It will be clear however that the use of the full asymptotic expansion (6.29) will also lead to full asymptotic expansions for the polynomials P_n, Q_n, and R_n, and for the remainder E_n.

7 Proofs of the asymptotic formulas

We now know the asymptotic behavior (6.28) of S as $n \to \infty$. We will trace back our steps to the original Riemann–Hilbert problem for Y to obtain asymptotics for the polynomials in the Hermite–Padé approximation problem for the exponential function.

7.1 Proof of Theorem 2.8

Proof. We start with the proof of the asymptotic formula (2.23) for P_n. Let K be a compact subset of $\mathbb{C} \setminus \Gamma_P$. We have the freedom to take the contours Γ_{P^-}, Γ_{P^+}, $\Gamma_{P^{++}}$ near Γ_P, and the circles Γ_1 and Γ_2 around z_1 and z_2 in such a way that K is in the exterior of these curves. Let $z \in K$. Then we follow the transformations $Y \mapsto U \mapsto T \mapsto S$. We see first from (5.3) that

$$P_n(z) = Y_{21}(z) = U_{21}(z) e^{-(n+1)\ell} e^{(n+1)g_P(z)}.$$

Then from the definition of T in (5.13) and (5.18), we get that $U_{21}(z) = T_{21}(z)$. We finally note that $T(z) = S(z)N(z)$, if z is outside Γ_3 and Γ_4, and $T(z) = S(z)M^{(j)}(z)$ if z is inside Γ_j with $j = 3, 4$. For $T_{21}(z)$, we get in either case

$$T_{21}(z) = S_{21}(z)N_{11}(z) + S_{22}(z)N_{21}(z) + S_{13}(z)N_{31}(z)$$

since the first columns of $M^{(3)}$ and $M^{(4)}$ agree with the first column of N.

Since $S = I + \mathcal{O}\left(\frac{1}{n}\right)$, and since N_{21} does not vanish on K, we get

$$P_n(z) = N_{21}(z)e^{-(n+1)\ell}e^{(n+1)g_P(z)}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right)$$

uniformly on K. Now we use the formula in (6.20) for N_{21} and we recall that $\ell = \log 2 - \pi i$, to obtain (2.23).

The proof of the asymptotic formula (2.24) for R_n is similar.

For Q_n we proceed differently, because of the way Q_n appears in the entries of Y. We take $z_0 \in \mathbb{C} \setminus \Gamma_Q$, and show that there is a neighborhood Δ of z_0 such that (2.25) holds uniformly for $z \in \Delta$. First we assume that z_0 belongs to the outside region D_∞. Then we can take the original contour Γ so that a neighborhood Δ of z_0 is in the outside region. Then for $z \in \Delta$, we have by (5.1)

$$Q_n(z) = z^{3n+2}Y_{22}(z) = U_{22}(z)z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))}.$$

We can open the contours around Γ_P and Γ_R so that Δ is in the exterior region to these contours. Then we have $U = T = SN = \left(I + \mathcal{O}\left(\frac{1}{n}\right)\right)N$, so that $U_{22}(z) = N_{22}(z)\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right)$ uniformly for $z \in \Delta$. This leads to the first formula in (2.25), if we use (4.24) and the formula for N_{22} in (6.20).

If $z_0 \in D_P$, then we can also open up the lenses around Γ_P and Γ_R so that a neighborhood Δ of z_0 is not contained in these lenses. Then we have for $z \in \Delta$, by (5.2)

$$Q_n(z) = E_n(z) - P_n(z)e^{-3nz} - R_n(z)e^{3nz}$$

and we need to find out, what is the dominant contribution as n gets large. We already have asymptotic formulas for P_n and R_n, from which it follows that

$$\frac{1}{n}\log|P_n(z)e^{-3nz}| \to \operatorname{Re}(g_P(z) - 3z - \ell), \tag{7.1}$$

$$\frac{1}{n}\log|R_n(z)e^{3nz}| \to \operatorname{Re}(g_R(z) + 3z - \ell). \tag{7.2}$$

For E_n it is easy to obtain in a similar way

$$\frac{1}{n}\log|E_n(z)| \to \operatorname{Re}(3\log z - g_P(z) - g_R(z)). \tag{7.3}$$

Now it turns out that for $z \in D_P$ the term $P_n(z)e^{-3nz}$ dominates. Indeed, we have by (4.12) and (4.13)

$$(g_P(z) - 3z - \ell) - (g_R(z) + 3z - \ell) = (g_P - g_R - 6z) = 2(-\varphi_P(z) + \varphi_R(z))$$

and we know that the real part of $-\varphi_P + \varphi_R$ is positive in D_P. We also have by (4.12)

$$(g_P(z) - 3z - \ell) - (3\log z - g_P(z) - g_R(z)) = 2g_P(z) + g_R(z) - 3\log z - 3z - \ell = -2\varphi_P(z)$$

and the real part of $-2\varphi_P$ is positive in D_P. In view of (7.1)–(7.3) we then obtain the formula for Q_n

$$Q_n(z) = -P_n(z)e^{-3nz}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right)$$

uniformly for $z \in \Delta$, and from what we already know about P_n,

$$Q_n(z) = -\frac{(-1)^{n+1}e^{ng_P(z)}}{2^n\sqrt{3\psi_P^4 + 1}}e^{-3nz}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right)$$

uniformly for $z \in \Delta$. This leads to the second line in formula (2.25) because $g_P(z) - 3z + \log(-\frac{1}{2}) = g_Q(z)$ for $z \in D_P$, see (4.9) and (4.23). The proof for the case $z_0 \in D_R$ is similar.

We finally have to consider the case that z_0 is on Γ_P or on Γ_R (but not one of the branch points). We consider only $z_0 \in \Gamma_P$, since $z_0 \in \Gamma_R$ will follow in the same way. So let $z_0 \in \Gamma_P \setminus \{z_1, z_2\}$. After opening up lenses around Γ_P we have the contour Γ_{P-} to the left of z_0, and the contour Γ_{P+} to the right. We can take a neighborhood Δ of z_0 that is strictly contained in the domain bounded by Γ_{P-} and Γ_{P+}. Then for $z \in \Delta \cap D_\infty$, we have

$$Q_n(z) = z^{3n+2}Y_{22}(z) = U_{22}(z)z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))}$$

and $U(z) = T(z)V_{P-}^{-1}(z)$. Now we have $U_{22}(z) = T_{22}(z)$, since the second column of V_{P-}^{-1} is simply $\begin{pmatrix}0 & 1 & 0\end{pmatrix}^T$. We open the circles Γ_1 and Γ_2 around z_1 and z_2 so that Δ is in the exterior. Then $T = SN = (I + \mathcal{O}(\frac{1}{n}))N$, and so

$$Q_n(z) = z^{3n+2}N_{22}(z)z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))}\left(I + \mathcal{O}\left(\frac{1}{n}\right)\right)$$

uniformly for $z \in \Delta \cap D_\infty$. Using (4.24) and the formula (6.20) for N_{22}, we then find (2.25).

For $z \in \Delta \cap D_P$, we have

$$\begin{aligned}
Q_n(z) &= E_n(z) - P_n(z)e^{-3nz} - R_n(z)e^{3nz} \\
&= z^{3n+2}Y_{22}(z) - e^{-3nz}Y_{21}z - e^{3nz}Y_{13}(z) \\
&= z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))}U_{22}(z) - e^{-3nz}e^{(n+1)(g_P(z)-\ell)}U_{21}(z) \\
&\quad -e^{3nz}e^{(n+1)(g_R(z)-\ell)}U_{13}(z).
\end{aligned}$$

Now we have that

$$U(z) = T(z)V_{P+}(z)V_{P++}(z)$$

with

$$V_{P+}(z)V_{P++}(z) = \begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)} & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix}.$$

Then

$$Q_n(z) = -e^{-3nz}e^{(n+1)(g_P(z)-\ell)}T_{21}(z)$$
$$+\left[z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))} - e^{-3nz}e^{(n+1)(g_P(z)-\ell)}z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)}\right]T_{22}(z)$$
$$+\left[z^{3n+2}e^{-(n+1)(g_P(z)+g_R(z))}ze^{-3z}e^{-2(n+1)\varphi_R(z)} - e^{3nz}e^{(n+1)(g_R(z)-\ell)}\right]T_{13}(z).$$

The funny thing is that the expressions multiplying $T_{22}(z)$ and $T_{13}(z)$ are exactly zero. This follows from (4.12) and (4.13). Thus only the first term remains. This gives

$$\begin{aligned} Q_n(z) &= -e^{g_P(z)-\ell}e^{ng_Q(z)}T_{21}(z) \\ &= -e^{g_P(z)-\ell}N_{21}(z)e^{ng_Q(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \\ &= \frac{1}{\sqrt{\psi_P^4(z)+1}}e^{ng_Q(z)}\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \end{aligned}$$

uniformly for $z \in \Delta \cap D_P$. In the last step we used (4.9) and the expression for N_{21}. This proves the asymptotic formula for $z \in \Delta \cap D_P$.

The asymptotic formula (2.26) for E_n is established in $D_P \cup D_R \cup \Gamma_Q$ again by following the transformations $Y \mapsto U \mapsto T \mapsto S$. The asymptotics in the other domains may be obtained by observing that the main contribution in the sum

$$E_n(z) = P_n(z)e^{-3nz} + Q_n(z) + R_n(z)e^{3nz}$$

is given by $P_n(z)e^{-3nz}$, $Q_n(z)$ or $R_n(z)e^{3nz}$ if z belongs to the domains $D_{\infty,P}$, $D_{\infty,U} \cup D_{\infty,L}$ or $D_{\infty,R}$ respectively. This completes the proof of Theorem 2.8. ☐

7.2 Proof of Theorem 2.5

Proof. The limits for the counting measures ν_{P_n}, ν_{Q_n}, and ν_{R_n} follow from the strong convergence results established in Theorem 2.8. The proof using the unicity theorem for logarithmic potentials (see e.g. [30, Theorem II.2.1]) is the same as the proof of [35, Theorem 2.1].

The proof for the limit of the measures ν_{E_n} is more difficult, since these measures have unbounded support and infinite mass. Our proof is different from the one given by Stahl [35].

Since E_n is an entire function of order 1 with a zero of multiplicity $3n + 2$ at the origin, we have by the Hadamard factorization theorem

$$E_n(z) = z^{3n+2}e^{A_n z + B_n}\prod_{j=1}^{\infty}\left(1 - \frac{z}{z_{j,n}}\right)e^{z/z_{j,n}},$$

with $A_n, B_n \in \mathbb{C}$ and where $z_{j,n}$, $j = 1, 2, \ldots$ are the zeros of E_n different from the origin. The zeros are counted according to their multiplicities, and we assume that they are arranged in non-decreasing absolute value, so that $0 < |z_{1,n}| \leq |z_{2,n}| \leq \cdots$. Taking logarithms, we get

$$\frac{1}{n}\log\left(z^{-3n-2}E_n(z)\right) = \frac{1}{n}(A_n z + B_n) + \int \left(\log\left(1 - \frac{z}{s}\right) + \frac{z}{s}\right) d\nu_{E_n}(s).$$

From the strong asymptotic formula (2.26), we obtain from this that

$$\lim_{n\to\infty}\left[\frac{1}{n}(A_n z + B_n) + \int \left(\log\left(1-\frac{z}{s}\right)+\frac{z}{s}\right) d\nu_{E_n}(s)\right] = g_E(z) \tag{7.4}$$

where

$$g_E(z) = \begin{cases} g_R(z) + 3z - 3\log z - \ell & \text{for } z \in D_{\infty,R}, \\ g_P(z) - 3z - 3\log z - \ell & \text{for } z \in D_{\infty,P}, \\ -g_P(z) - g_R(z) & \text{for } z \in \mathbb{C} \setminus (\overline{D_{\infty,R} \cup D_{\infty,P}}). \end{cases} \tag{7.5}$$

Note that g_E is analytic across Γ_P and Γ_R because of the relations (4.14)–(4.15), so that g_E is analytic in $\mathbb{C} \setminus \Gamma_E$. The limit (7.4) holds uniformly for z in compact subsets of $\mathbb{C} \setminus \Gamma_E$.

We first show that

$$g_E(z) = \int \left(\log\left(1-\frac{z}{s}\right)+\frac{z}{s}\right) d\mu_E(s) - g_P(0) - g_R(0) \tag{7.6}$$

for $z \in \mathbb{C} \setminus \Gamma_E$. To prove (7.6) we assume $z \in \mathbb{C} \setminus (\overline{D_{\infty,R} \cup D_{\infty,P}})$. We take the derivative of the right-hand side of (7.6), which is

$$\begin{aligned} \int \left(\frac{1}{z-s}+\frac{1}{s}\right) d\mu_E(s) &= \frac{3}{2\pi i}\int_{\Gamma_{E,1}\cup\Gamma_{E,2}} \left(\frac{1}{z-s}+\frac{1}{s}\right)(\psi_Q-\psi_P)(s)ds \\ &\quad + \frac{3}{2\pi i}\int_{\Gamma_{E,3}\cup\Gamma_{E,4}} \left(\frac{1}{z-s}+\frac{1}{s}\right)(\psi_Q-\psi_R)(s)ds. \end{aligned} \tag{7.7}$$

The integral over $\Gamma_{E,1} \cup \Gamma_{E,2}$ can be deformed to the integral over the −side of Γ_P, which leads to

$$\begin{aligned} \frac{3}{2\pi i}\int_{\Gamma_{E,1}\cup\Gamma_{E,2}} \left(\frac{1}{z-s}+\frac{1}{s}\right)(\psi_Q-\psi_P)(s)ds &= \frac{3}{2\pi i}\int_{\Gamma_P} \left(\frac{1}{z-s}+\frac{1}{s}\right)(\psi_Q-\psi_P)_-(s)ds \\ &= \frac{3}{2\pi i}\oint_{\gamma} \left(\frac{1}{z-s}+\frac{1}{s}\right)\psi_P(s)ds \end{aligned}$$

where γ is a closed contour going around Γ_P in the negative direction and with z and 0 outside γ, see also the proof of Lemma 4.3. We compute the integral using the residue theorem for the exterior of γ, for which there is a contribution from the poles at z and 0. The result is that the first term on the right-hand side of (7.7) is equal to $-3\psi_P(z) + 3\psi_P(0)$. In the same way, we find that the second term is equal to $-3\psi_R(z) + 3\psi_R(0)$. Using (4.4), (4.5) and the fact that $\psi_R(0) = \sqrt{1/3} = -\psi_P(0)$, we see that the derivative of the right-hand side of (7.6) is equal to $-g_P'(z) - g_R'(z)$ for $z \in \mathbb{C} \setminus (\overline{D_{\infty,R} \cup D_{\infty,P}})$. Because of the definition of g_E in (7.5) we then get (7.6) in case $z \in \mathbb{C} \setminus (\overline{D_{\infty,R} \cup D_{\infty,P}})$. The formula (7.6) in the other domains follows by analytical continuation.

Knowing (7.6) we get from (7.4) and (7.5) that $\lim_{n\to\infty} \frac{1}{n}A_n = 0$, $\lim_{n\to\infty} \frac{1}{n}B_n = -g_P(0) - g_R(0)$, and

$$\lim_{n\to\infty}\int \left(\log\left(1-\frac{z}{s}\right)+\frac{z}{s}\right) d\nu_{E_n}(s) = \int \left(\log\left(1-\frac{z}{s}\right)+\frac{z}{s}\right) d\mu_E(s) \tag{7.8}$$

for $z \in \mathbb{C} \setminus \Gamma_E$.

We next consider the measures ρ_n defined by

$$d\rho_n(s) = \frac{1}{|s|^2} d\nu_{E_n}(s).$$

It will be our goal to prove that

$$\sup_{n\in\mathbb{N}} \int d\rho_n < +\infty \tag{7.9}$$

and that the sequence (ρ_n) is tight, that is,

$$\forall \varepsilon > 0 : \exists R > 0 : \forall n \in \mathbb{N} : \int_{|s|\geq R} d\rho_n(s) \leq \varepsilon, \tag{7.10}$$

see [6, Chapter 6]. We obtain (7.9) and (7.10) from lower bounds on the absolute values of the zeros $z_{j,n}$ of E_n.

Let $N_n(r)$ be the number of non-zero roots of E_n with absolute value $\leq r$. From the theory of entire functions, we have that

$$N_n(r) < \log \max_{|z|=er} |f_n(z)| - \log |f_n(0)|, \tag{7.11}$$

see [25, Section 2.5], where

$$f_n(z) = z^{-3n-2} E_n(z). \tag{7.12}$$

The integral representations (1.4)–(1.6) and (1.7) imply that

$$E_n(z) = \frac{(-1)^{n+1} n!}{(3n)^n} \frac{1}{2\pi i} \oint_{C_R} \frac{e^{3nzw} dw}{[w(w^2-1)]^{n+1}}, \tag{7.13}$$

where C_R is a circle centered at the origin with radius $R > 1$. From (7.13) it immediately follows that

$$E_n(z) = \frac{(-1)^{n+1} n! (3n)^{2n+2}}{(3n+2)!} z^{3n+2} + \mathcal{O}(z^{3n+3}) \qquad \text{as } z \to 0,$$

and so

$$f_n(0) = \frac{(-1)^{n+1} n! (3n)^{2n+2}}{(3n+2)!}. \tag{7.14}$$

Using $E_n(z) = P_n(z)e^{-3nz} + Q_n(z) + R_n(z)e^{3nz}$ and the fact that the polynomials P_n, Q_n, and R_n, of leading coefficients $(-1/2)^{n+1}$, 1, and $(-1/2)^{n+1}$ respectively, have their zeros in a compact set (independent of n), we easily get that $|E_n(z)| \leq e^{4n|z|}$ for every $n \in \mathbb{N}$ and for every $|z| \geq R$ with R sufficiently large, say $R \geq R_0 \geq 1$. Then, from (7.11), (7.12) and (7.14) we see that there exists a constant $C > 0$ so that $N_n(r) < Cnr$ if $r > R_0$. Hence, for every j with $|z_{j,n}| > R_0$, we have $j \leq N_n(|z_{j,n}|) < Cn|z_{j,n}|$, so that

$$|z_{j,n}| > \frac{j}{Cn} \qquad \text{if } |z_{j,n}| > R_0. \tag{7.15}$$

This is our required estimate on $|z_{j,n}|$.

To estimate

$$\int d\rho_n = \frac{1}{n} \sum_{j=1}^{\infty} \frac{1}{|z_{j,n}|^2},$$

we note that there are at most CnR_0 zeros with absolute value $\leq R_0$. So for $j > CnR_0$, we have $|z_{j,n}| > R_0$, and we can use the estimate (7.15). For $j \leq CnR_0$, we use the fact that the zeros can only accumulate on Γ_E, which implies that there is a $\delta > 0$ so that $|z_{j,n}| > \delta$ for every j and n. Then for every $n \in \mathbb{N}$,

$$\int d\rho_n \leq \frac{1}{n}\sum_{j=1}^{[CnR_0]} \frac{1}{\delta^2} + \frac{1}{n}\sum_{j=[CnR_0]+1}^{\infty} \frac{(Cn)^2}{j^2} \leq \frac{CR_0}{\delta^2} + \frac{C^2 n}{[CnR_0]},$$

and (7.9) follows.

Next, we estimate for $R > R_0$,

$$\begin{aligned} \int_{|s|\geq R} d\rho_n(s) &= \frac{1}{n}\sum_{j=N_n(R)+1}^{\infty} \frac{1}{|z_{j,n}|^2} \leq \frac{1}{n}\sum_{j=N_n(R)+1}^{\infty} \min\left(\frac{1}{R^2}, \frac{C^2 n^2}{j^2}\right) \\ &\leq \frac{1}{n}\sum_{j=1}^{[CnR]} \frac{1}{R^2} + \frac{1}{n}\sum_{j=[CnR]+1}^{\infty} \frac{C^2 n^2}{j^2} \leq \frac{C}{R} + \frac{C^2 n}{[CnR]}, \end{aligned}$$

and this shows that, given $\varepsilon > 0$, we can choose $R > R_0$ sufficiently large so that $\int_{|s|\geq R} d\rho_n(s) \leq \varepsilon$ for every n. Hence (7.10) holds and the sequence (ρ_n) is tight.

Now let ρ^* be the limit of a weakly convergent subsequence of (ρ_n), where $\rho_n \to \rho$ weakly means that $\int f d\rho_n \to \int f d\rho$ for every bounded continuous function f. Because of (7.9), we have that $\int d\rho^* < \infty$. Let $d\nu^*(s) = |s|^2 d\rho^*(s)$. Then (7.8) implies

$$\int \left(\log\left(1 - \frac{z}{s}\right) + \frac{z}{s}\right) d\nu^*(s) = \int \left(\log\left(1 - \frac{z}{s}\right) + \frac{z}{s}\right) d\mu_E(s) \tag{7.16}$$

for every $z \in \mathbb{C} \setminus \Gamma_E$. Let $R > 0$ and let ν_R^* and $\mu_{E,R}$ be the restrictions to $\{|z| < R\}$ of ν^* and μ_E, respectively. Then we get by taking the real part and rewriting (7.16) that

$$\int \log|z - s| d\nu_R^*(s) = \int \log|z - s| d\mu_{E,R}(s) + h_R(z),$$

where h_R is harmonic in $\{|z| < R\}$. Since Γ_E has two–dimensional Lebesgue measure 0, the unicity theorem [30, Theorem II.2.1] applies, showing that ν_R^* and $\mu_{E,R}$ are equal. As R is arbitrary, we have that $\nu^* = \mu_E$ and so $d\rho^*(s) = \frac{1}{|s|^2} d\mu_E(s)$. Since the sequence (ρ_n) is tight, and $\frac{1}{|s|^2} d\mu_E(s)$ is the only possible limit of a weakly convergent subsequence, the full sequence (ρ_n) converges weakly to $\frac{1}{|s|^2} d\mu_E(s)$, see [6, Chapter 6]. Then it also follows that the measures ν_{E_n} converge to μ_E in the sense indicated in the theorem.

This completes the proof of the theorem. □

7.3 Proof of Theorem 2.9

Proof. Away from Γ_Q we have that one of the terms in (2.27) dominates the others, and then (2.27) reduces to (2.25). So it suffices to prove that (2.27) holds uniformly for z in a neighborhood Δ of a point $z_0 \in \Gamma_Q \setminus \{z_1, z_2, z_3, z_4\}$. So let $z_0 \in \Gamma_Q \setminus \{z_1, z_2, z_3, z_4\}$. We take the contours $\Gamma_{P^{++}}$ and $\Gamma_{R^{++}}$ so that we can choose a disk Δ centered at z_0 that lies in the region bounded by $\Gamma_{P^{++}}$, $\Gamma_{R^{++}}$, Γ_{U^m} and Γ_{L^m}.

From (1.3) we know that

$$Q_n(z) = E_n(z) - P_n(z)e^{-3nz} - R_n(z)e^{3nz}. \tag{7.17}$$

We already know asymptotic expressions for the polynomials P_n, R_n and the remainder function E_n that are valid in Δ, namely (2.23), (2.24) and the last equation in (2.26). Plugging these three formulas into (7.17), we deduce

$$\begin{aligned} Q_n(z) = z^{3n}e^{-n[g_P(z)+g_R(z)]} \Bigg[& \frac{e^{n[2g_P(z)+g_R(z)-\ell-3\log z-3z]}}{\sqrt{3\psi_P^4(z)+1}} \left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) - \frac{\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)}{\sqrt{3\psi_Q^4(z)+1}} \\ & + \frac{e^{n[g_P(z)+2g_R(z)-\ell-3\log z+3z]}}{\sqrt{3\psi_R^4(z)+1}} \left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\Bigg]. \end{aligned}$$

Using (4.12) and (4.13) we are led to (2.27), uniformly for $z \in \Delta$.

The proof of (2.28) is similar. We use (2.23), (2.24) and the first equation in (2.25). ☐

7.4 Proof of Theorem 2.10

Proof. Away from Γ_P one of the terms in (2.29) is dominating the other, and then (2.29) leads to (2.23). So it suffices to consider $z_0 \in \Gamma_P \setminus \{z_1, z_2\}$, and prove that (2.29) holds uniformly for z in a disk Δ centered at z_0. We may choose this disk so that it is disjoint from Γ_1 and Γ_2, and also so that it is contained in the region bounded by Γ_{P-} and Γ_{P+}, see Figure 13.

We consider $z \in \Delta$. Then we have as before $P_n(z) = U_{21}(z)e^{(n+1)(g_P(z)-\ell)}$, but the relation between T and U is $U(z) = T(z)V_{P-}^{-1}(z)$, or $U(z) = T(z)V_{P+}(z)V_{P++}$ depending on whether z lies in the region bounded by Γ_P and Γ_{P-} (the $-$side of Γ_P) or in the region bounded by Γ_P and Γ_{P+} (the $+$side). This gives

$$U_{21}(z) = T_{21}(z) \pm z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)}T_{22}(z),$$

for z on the $\pm$side of Γ_P.

From the fact that $T = SN$ with $S = I + \mathcal{O}(\frac{1}{n})$, we get

$$\begin{aligned} P_n(z) = e^{(n+1)(g_P(z)-\ell)} & \left(N_{21}(z) \pm z^{-1}e^{2(n+1)\varphi_P(z)-3z}N_{22}(z)\right)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \\ & + \left(N_{11}(z) \pm z^{-1}e^{2(n+1)\varphi_P(z)3z}N_{12}(z)\right)\mathcal{O}\left(\frac{1}{n}\right) \\ & + \left(N_{31}(z) \pm z^{-1}e^{2(n+1)\varphi_{P-}(z)3z}N_{3,2}(z)\right)\mathcal{O}\left(\frac{1}{n}\right). \end{aligned} \tag{7.18}$$

Since all functions N_{jk} do not vanish in Δ, and since $e^{\ell} = -2$, we obtain

$$(-2)^{n+1}P_n(z) = e^{(n+1)g_P(z)}\left[N_{21}(z)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right) \pm z^{-1}e^{2(n+1)\varphi_P(z)3z}N_{22}(z)\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\right].$$

Now we use (4.12) and the formulas in (6.20) for N_{21} and N_{22}, and (2.29) follows.

The proof of the asymptotic formula (2.30) for R_n is similar. ☐

7.5 Proof of Theorem 2.11

Proof. If we unravel all the transformations for $z \in \Delta_1$, we find that $Y(z)$ is a product of no less than 10 to 14 matrices, the exact number depending on the region where z is. Recall that z belongs to one of the four regions $f^{-1}(I)$, $f^{-1}(II)$, $f^{-1}(III)$, and $f^{-1}(IV)$, see Figure 12. For $z \in f_1^{-1}(III)$, we have the following, where $s = (n+1)^{2/3} f_1(z)$,

$$\begin{aligned}
Y(z) = e^{i\pi/6}\sqrt{\pi} &\begin{pmatrix} e^{(n+1)\ell/3} & 0 & 0 \\ 0 & e^{-2(n+1)\ell/3} & 0 \\ 0 & 0 & e^{(n+1)\ell/3} \end{pmatrix} S(z)N(z) \\
&\begin{pmatrix} z^{1/2}e^{3z/2} & 0 & 0 \\ 0 & z^{-1/2}e^{-3z/2} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ i & i & 0 \\ 0 & 0 & 2e^{-i\pi/6} \end{pmatrix} \begin{pmatrix} s^{1/4} & 0 & 0 \\ 0 & s^{-1/4} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \operatorname{Ai}(s) & -\operatorname{Ai}(\omega_3^2 s) & 0 \\ \operatorname{Ai}'(s) & -\omega_3^2 \operatorname{Ai}'(\omega_3^2 s) & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
&\begin{pmatrix} e^{-i\pi/6} & 0 & 0 \\ 0 & e^{i\pi/6} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} z^{-1/2}e^{-3z/2}e^{(n+1)\varphi_P(z)} & 0 & 0 \\ 0 & z^{1/2}e^{3z/2}e^{-(n+1)\varphi_P(z)} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
&\begin{pmatrix} 1 & 0 & 0 \\ z^{-1}e^{-3z}e^{2(n+1)\varphi_P(z)} & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & ze^{-3z}e^{-2(n+1)\varphi_R(z)} & 1 \end{pmatrix} \\
&\begin{pmatrix} e^{-(n+1)\ell/3} & 0 & 0 \\ 0 & e^{2(n+1)\ell/3} & 0 \\ 0 & 0 & e^{-(n+1)\ell/3} \end{pmatrix} \begin{pmatrix} e^{(n+1)g_P(z)} & 0 & 0 \\ 0 & e^{-(n+1)[g_P(z)+g_R(z)]} & 0 \\ 0 & 0 & e^{(n+1)g_R(z)} \end{pmatrix}.
\end{aligned} \tag{7.19}$$

Recall that $P_n(z)$ is the $(2,1)$ entry of $Y(z)$. We multiply (7.19) with the first unit vector $\begin{pmatrix} 1 & 0 & 0 \end{pmatrix}^T$, and we obtain after some simple calculations, where $*$ denotes an unspecified unimportant entry

$$\begin{pmatrix} * \\ P_n(z) \\ * \end{pmatrix} = \frac{\sqrt{\pi} e^{(n+1)(g_P(z)+\varphi_P(z))}}{(-2)^{n+1}} S(z)N(z) \begin{pmatrix} 1 & 0 & 0 \\ 0 & z^{-1}e^{-3z} & 0 \\ 0 & 0 & * \end{pmatrix} \begin{pmatrix} 1 & -1 & 0 \\ i & i & 0 \\ 0 & 0 & * \end{pmatrix} \begin{pmatrix} s^{1/4}\operatorname{Ai}(s) \\ s^{-1/4}\operatorname{Ai}'(s) \\ 0 \end{pmatrix}. \tag{7.20}$$

Similar calculations for z in the other regions near z_1 lead to exactly the same expression (7.20). Then further calculations show that for $z \in \Delta_1$,

$$\begin{aligned}
(-2)^{n+1} P_n(z) = \sqrt{\pi} &e^{(n+1)(g_P(z)+\varphi_P(z))} \\
&\left[\sum_{k=1}^{3} S_{2k}(z) \left(N_{k1}(z) + iz^{-1}e^{-3z} N_{k2}(z) \right) (n+1)^{1/6} f_1(z)^{1/4} \operatorname{Ai}((n+1)^{2/3} f_1(z)) \right. \\
&\left. + \sum_{k=1}^{3} S_{2k}(z) \left(-N_{k1}(z) + iz^{-1}e^{-3z} N_{k2}(z) \right) (n+1)^{-1/6} f_1(z)^{-1/4} \operatorname{Ai}'((n+1)^{2/3} f_1(z)) \right].
\end{aligned} \tag{7.21}$$

From the jump condition (6.1) for N on Γ_P it easily follows that for each k,

$$\left(N_{k1}(z) + iz^{-1}e^{-3z} N_{k2}(z) \right)_+ = i \left(N_{k1}(z) + iz^{-1}e^{-3z} N_{k2}(z) \right)_-, \qquad z \in \Gamma_P.$$

The fourth root in $f_1(z)^{1/4}$ is defined with a cut along Γ_P, and on Γ_P there is a jump

$$\left(f_1(z)^{1/4}\right)_+ = -i\left(f_1(z)^{1/4}\right)_-, \qquad z \in \Gamma_P.$$

Thus the products

$$\left(N_{k1}(z) + iz^{-1}e^{-3z}N_{k2}(z)\right) f_1(z)^{1/4}$$

are analytic across Γ_P. Since the functions N_{kj} have a fourth root singularity at z_1, and $f_1(z)^{1/4}$ has a fourth root zero at z_1, these functions are non-zero at z_1, and therefore also in a neighborhood of z_1. By shrinking Δ_1, if necessary, we may assume that these functions are without zeros in Δ_1. Similarly, we have that

$$\left(-N_{k1}(z) + iz^{-1}e^{-3z}N_{k2}(z)\right) f_1(z)^{-1/4}$$

is analytic and without zeros in Δ_1.

Now we recall that $S(z) = I + \mathcal{O}\left(\frac{1}{n}\right)$. So it follows that in the two sums in (7.21) the terms with $k = 2$ are dominating, and we get

$$\begin{aligned}(-2)^{n+1}P_n(z) = {}& \sqrt{\pi}e^{(n+1)(g_P(z)+\varphi_P(z))} \\ & \left[n^{1/6}\left(N_{21}(z) + iz^{-1}e^{-3z}N_{22}(z)\right) f_1(z)^{1/4}\operatorname{Ai}((n+1)^{2/3}f_1(z))\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\right. \\ & \left. +n^{-1/6}\left(-N_{21}(z) + iz^{-1}e^{-3z}N_{22}(z)\right) f_1(z)^{-1/4}\operatorname{Ai}'((n+1)^{2/3}f_1(z))\left(1+\mathcal{O}\left(\frac{1}{n}\right)\right)\right]. \end{aligned} \tag{7.22}$$

This proves (2.34). We have also shown above that the functions h_1 and h_2 are analytic and without zeros in Δ_1.

For the asymptotic formula for Q_n near z_1, we proceed in a similar way. Actually, it is easier to do the calculations for z in the region $f_1^{-1}(II)$ or $f_1^{-1}(I)$. The result is that for $z \in \Delta_1$,

$$\begin{aligned}\begin{pmatrix} * \\ Q_n(z) \\ * \end{pmatrix} = {}& \sqrt{\pi}e^{3z}e^{-(n+1)(g_P(z)+\varphi_P(z)+g_R(z)-3\log z)}S(z)N(z) \\ & \begin{pmatrix} 1 & 0 & 0 \\ 0 & z^{-1}e^{-3z} & 0 \\ 0 & 0 & * \end{pmatrix}\begin{pmatrix} 1 & -1 & 0 \\ i & i & 0 \\ 0 & 0 & * \end{pmatrix}\begin{pmatrix} s^{1/4}\omega_3^2\operatorname{Ai}(\omega_3^2 s) \\ s^{-1/4}\omega_3\operatorname{Ai}'(\omega_3^2 s) \\ 0 \end{pmatrix}, \end{aligned} \tag{7.23}$$

where again $s = (n+1)^{2/3}f_1(z)$. We make use of (4.12) to replace the exponential factor on the right-hand side of (7.23) by $e^{(n+1)(g_P(z)+\varphi_P(z)-3z-l)}$. Then, we obtain (2.35) from (7.23) in the same way as we obtained (2.34) from (7.20).

For the asymptotic formula for E_n near z_1, we may again proceed in a similar way, doing the calculations for z in the region $f_1^{-1}(III)$ or $f_1^{-1}(IV)$. □

7.6 Proof of Corollary 2.12

Proof. The behavior of the extreme zeros of P_n near z_1 follows from the asymptotic formula (2.34). Indeed, we consider the function

$$F_n(t) = \frac{(-2)^{n+1}e^{-(n+1)(g_P(z)+\varphi_P(z))}}{\sqrt{\pi}n^{1/6}h_1(z)}P_n(z), \qquad \text{where } z = z_1 + tn^{-2/3}.$$

Then F_n has zeros $t_{\nu,n} = (z^P_{\nu,n} - z_1)n^{2/3}$, $\nu = 1, 2, \ldots$, and these zeros are ordered by increasing absolute value. Because of (2.34) we have that

$$F_n(t) = \mathrm{Ai}\left((n+1)^{2/3} f_1(z_1 + tn^{-2/3})\right)\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) + n^{-1/3}\frac{h_2(z_1 + tn^{-2/3})}{h_1(z_1 + tn^{-2/3})}\,\mathrm{Ai}'\left((n+1)^{2/3} f_1(z_1 + tn^{-2/3})\right)\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right). \quad (7.24)$$

Since f_1 is an analytic function with a simple zero at z_1 and $f_1'(z_1) = c_1$ (see (2.32), (2.33)), and since h_1 and h_2 are nonzero analytic functions near z_1, we get by expanding these functions near z_1 that

$$F_n(t) = \mathrm{Ai}\left(tc_1 + \mathcal{O}\left(n^{-2/3}\right)\right) + n^{-1/3}\frac{h_2(z_1)}{h_1(z_1)}\,\mathrm{Ai}'\left(tc_1 + \mathcal{O}\left(n^{-2/3}\right)\right) + \mathcal{O}\left(n^{-2/3}\right). \quad (7.25)$$

Now, expanding the Airy function Ai near tc_1, and observing that the second term of the sum in the right-hand side of (7.25) is of order $n^{-1/3}$, we get simply

$$F_n(t) = \mathrm{Ai}(tc_1) + \mathcal{O}\left(n^{-1/3}\right). \quad (7.26)$$

The $\mathcal{O}$-term holds uniformly on compact subsets of the complex t-plane. From Hurwitz' theorem it follows that for every fixed $\nu \in \mathbb{N}$, we have

$$\lim_{n\to\infty} t_{\nu,n} = -\frac{\iota_\nu}{c_1}.$$

Using the fact that $-\iota_\nu$ is a simple zero of the Airy function, we obtain from (7.26) that

$$t_{\nu,n} = -\frac{\iota_\nu}{c_1} + \mathcal{O}\left(n^{-1/3}\right).$$

This proves (2.40), since $z^P_{\nu,n} = z_1 + t_{\nu,n}n^{-2/3}$.

The formulas (2.41) and (2.42) for the extreme zeros of Q_n and E_n near z_1 are obtained in a similar way from the asymptotics of Q_n and E_n near z_1, given by (2.35) and (2.36), respectively. □

7.7 Proof of Theorem 2.13

Proof. We consider z in a compact subset K of D_R. By the third equation in (2.25), we have

$$\lim_{n\to\infty} \frac{1}{n}\log|Q_n^2(z)| = 2\mathrm{Re}\, g_Q(z),$$

and by (2.23) and (2.24),

$$\lim_{n\to\infty} \frac{1}{n}\log|P_n(z)R_n(z)| = \mathrm{Re}\,(g_P(z) + g_R(z) - 2\ell).$$

In D_R we may use (4.26), so that

$$2\mathrm{Re}\, g_Q(z) - \mathrm{Re}\,(g_P(z) + g_R(z) - 2\ell) = \mathrm{Re}\,(g_R(z) - g_P(z) + 6z).$$

By (4.12) and (4.13), this is $2\mathrm{Re}\,(\varphi_P - \varphi_R)$ which is positive in D_R. Thus, $4P_nR_n/Q_n^2$ is exponentially small, and there is a choice of the square root so that

$$\sqrt{Q_n^2(z) - 4P_n(z)R_n(z)} = Q_n(z)\left(1 + \mathcal{O}(e^{-cn})\right),$$

as $n \to \infty$, uniformly for $z \in K$, for some constant $c > 0$. Then we choose the $-$ sign in (2.44) so that

$$X_n(z) = \frac{-Q_n(z) - \sqrt{Q_n^2(z) - 4P_n(z)R_n(z)}}{2R_n(z)} = -\frac{Q_n(z)}{R_n(z)}\left(1 + \mathcal{O}(e^{-cn})\right).$$

Now, we use (2.24) and the third equation in (2.25) to obtain

$$X_n(z) = \frac{-(-2)^{n+1}e^{ng_Q(z)}}{2e^{ng_R(z)}}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right) = e^{3nz}\left(1 + \mathcal{O}\left(\frac{1}{n}\right)\right),$$

where the last equality uses (4.26). This proves (2.45) for $z \in D_R$.

The case $z \in D_P$ is similar, but now we have to take the $+$ sign in the formula (2.44) for X_n. This completes the proof of Theorem 2.13. $\square$

References

[1] A.I. Aptekarev, *Convergence of rational approximations to a set of exponential functions.* (Russian) Vestnik Moskov. Univ. Ser. I Mat. Mekh. 1981, no. 1, 68–74, 108. English translation: Moscow Univ. Math. Bull. 36 (1981), no. 1, 81–86.

[2] A.I. Aptekarev and H. Stahl, *Asymptotics of Hermite–Padé polynomials*, *in* “Progress in Approximation Theory” (A.A. Gonchar and E.B. Saff, Eds.), 127–167, Springer Verlag, 1992.

[3] J. Baik, P. Deift, K. McLaughlin, P. Miller, and X. Zhou, *Optimal tail estimates for directed last passage site percolation with geometric random variables*, Adv. Theor. Math. Phys. **5** (2001), 1207–1250.

[4] J. Baik, T. Kriecherbauer, K. T.-R. McLaughlin, and P.D. Miller, *Uniform asymptotics for polynomials orthogonal with respect to a general class of discrete weights and universality results for associated ensembles: announcement of results*, preprint, Internat. Math. Res. Notices **2003** (2003), 821–858.

[5] G.A. Baker and D.S. Lubinsky, *Convergence theorems for rows of differential and algebraic Hermite–Padé approximations*, J. Comput. Appl. Math. **18** (1987), 29–52.

[6] P. Billingsley, *Convergence of Probability Measures*, John Wiley & Sons, New York, 1968.

[7] P. Bleher and A. Its, *Semiclassical asymptotics of orthogonal polynomials, Riemann–Hilbert problem, and the universality in the matrix model*, Ann. of Math. **150** (1999), 185–266.

[8] P. Bleher and A. Its, *Double scaling limit in the random matrix model: The Riemann-Hilbert approach* Comm. Pure Appl. Math. **56** (2003), 433–516.

[9] M. de Bruin, *Some aspects of simultaneous rational approximation*, Numerical analysis and mathematical modelling, Banach Cent. Publ. 24, 1990, pp. 51–84.

[10] P.B. Borwein, *Quadratic Hermite–Padé approximation to the exponential function*, Constr. Approx. **2** (1986), 291–302.

[11] G.V. Chudnovsky, *Hermite–Padé approximations to exponential functions and elementary estimates of the measure of irrationality of π*, Lecture Notes in Math. vol. 925, pp. 299–322, Springer Verlag, New York, 1982.

[12] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann–Hilbert Approach*, Courant Lecture Notes 3, New York University, 1999; Amer. Math. Soc., Providence RI, 2000.

[13] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, *Uniform asymptotics of polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory*, Commun. Pure Appl. Math. **52** (1999), 1335–1425.

[14] P. Deift, T. Kriecherbauer, K.T-R McLaughlin, S. Venakides, and X. Zhou, *Strong asymptotics of orthogonal polynomials with respect to exponential weights*, Commun. Pure Appl. Math. **52** (1999), 1491–1552.

[15] P. Deift and X. Zhou, *A steepest descent method for oscillatory Riemann–Hilbert problems: asymptotics for the MKdV equation*, Ann. of Math. **137** (1993), 295–368.

[16] K.A. Driver, *Non-diagonal quadratic Hermite–Padé approximation to the exponential function*, J. Comp. Appl. Math. **65** (1995), 125–134.

[17] K.A. Driver and N.M. Temme, *On polynomials related with Hermite–Padé approximations to the exponential function*, J. Approx. Theory **95** (1998), 101–122.

[18] A.S. Fokas, A.R. Its, and A.V. Kitaev, *The isomonodromy approach to matrix models in 2D quantum gravity*, Commun. Math. Phys. **147** (1992), 395–430.

[19] C. Hermite, *Sur la fonction exponentielle*, C. R. Acad. Sci. Paris **77** (1873), 18–24, 74–79, 226–233.

[20] T. Kriecherbauer and K.T-R McLaughlin, *Strong asymptotics of polynomials orthogonal with respect to Freud weights*, Internat. Math. Res. Notices **1999** (1999), 299–333.

[21] A.B.J. Kuijlaars, *Riemann–Hilbert analysis for orthogonal polynomials*, Lecture notes for Summer School in Orthogonal Polynomials and Special Functions, manuscript 2002.

[22] A.B.J. Kuijlaars and K.T-R McLaughlin, *Riemann–Hilbert analysis for Laguerre polynomials with large negative parameter*, Computational Methods and Function Theory **1** (2001), 205–233.

[23] A.B.J. Kuijlaars and K.T-R McLaughlin, *Asymptotic zero behavior of Laguerre polynomial with negative parameter*, math.CA/0205175, to appear in Constr. Approx.

[24] A.B.J. Kuijlaars, K.T-R McLaughlin, W. Van Assche, and M. Vanlessen, *The Riemann–Hilbert approach to strong asymptotics for orthogonal polynomials on* $[-1,1]$, preprint math.CA/0111252.

[25] B.Y. Levin, *Lectures on entire functions*, Transl. Math. Monographs Vol. 150, Amer. Math. Soc., Providence R.I., 1996.

[26] K. Mahler, *Applications of some formulas by Hermite to the approximation of exponentials and logarithms*, Math. Ann. **168** (1967), 200–227.

[27] J. Nuttall, *Asymptotics of diagonal Hermite–Padé approximants*, J. Approx. Theory **42** (1984), 299–386.

[28] F. Olver, *Asymptotics and Special Functions*, Academic Press, San Diego, 1974.

[29] C. Pommerenke, *Univalent Functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.

[30] E.B. Saff and V. Totik, *Logarithmic Potentials with External Fields*, Springer-Verlag, Berlin, New York, 1997.

[31] E.B. Saff and R.S. Varga, *On the zeros and poles of Padé approximants to* e^z, Numer. Math. **25** (1975), 1–14.

[32] E.B. Saff and R.S. Varga, *On the zeros and poles of Padé approximants to* e^z. II., Padé and rational approximations: theory and applications (E.B. Saff, R.S. Varga, eds.), pp. 195–213. New York, Academic Press, 1977.

[33] E.B. Saff and R.S. Varga, *On the zeros and poles of Padé approximants to* e^z. III., Numer. Math., Vol. **30** (1978), 241–266.

[34] H. Stahl, *Asymptotics for quadratic Hermite–Padé polynomials associated with the exponential function*, Electronic Trans. Num. Anal. **14** (2002), 193–220.

[35] H. Stahl, *Quadratic Hermite–Padé polynomials associated with the exponential function*, manuscript.

[36] H. Stahl, *Asymptotic distributions of zeros of quadratic Hermite–Padé polynomials associated with the exponential function*, manuscript.

[37] K. Strebel, Quadratic Differentials, Springer-Verlag, Berlin, 1984.

[38] G. Szegő, *Über eine Eigenschaft der Exponentialreihe*, Sitzungsber. Berl. Math. Ges. **23**, 50–64, 1924.

[39] W. Van Assche, J. S. Geronimo, and A. B. J. Kuijlaars, *Riemann–Hilbert problems for multiple orthogonal polynomials*, in 'Special Functions 2000: Current Perspective and Future Directions' (J. Bustoz et al., eds.), NATO Science Series II. Mathematics, Physics and Chemistry Vol. 30, Kluwer, Dordrecht, 2001, pp. 23–59.

[40] R.S. Varga and A.J. Carpenter, *Asymptotics for the zeros and poles of normalized Padé approximants to* e^z, Numer. Math. **68** (1994), 169–185.

[41] F. Wielonsky, *Asymptotics of diagonal Hermite–Padé approximants to e^z*, J. Approx. Theory **90** (1997), 283–298.

[42] F. Wielonsky, *Some properties of Hermite–Padé approximants to e^z*, in 'Continued Fractions: From Analytic Number Theory to Constructive Approximation' (B. C. Berndt, F. Gesztesy, eds.), Contemporary Mathematics **236**, Amer. Math. Soc., Providence RI, 1999, 369–379.

`A. B. J. Kuijlaars (arno@wis.kuleuven.ac.be)`
`W. Van Assche (walter@wis.kuleuven.ac.be)`
`Department of Mathematics`
`Katholieke Universiteit Leuven`
`Celestijnenlaan 200B`
`B-3001 Leuven, BELGIUM`

`F. Wielonsky (wielonsk@ano.univ-lille1.fr)`
`Laboratoire de Mathématiques appliquées`
`FRE CNRS 2222 - Bat. M2`
`Université des Sciences et Technologies Lille 1`
`F-59655 Villeneuve d'Ascq Cedex, FRANCE`